arXiv:0912.1024v1 [math.RA] 5 Dec 2009

SYMMETRIZABLE INTERSECTION MATRICES AND THEIR ROOT SYSTEMS

LIANGANG PENG AND MANG XU

Abstract. In this paper we study symmetrizable intersection matrices, namely generalized intersection matrices introduced by P. Slodowy such that they are symmetrizable. Every such matrix can be naturally associated with a root basis and a Weyl root system. Using d-fold affinization matrices we give a classification, up to braid-equivalence, for all positive semi-definite symmetrizable intersection matrices. We also give an explicit structure of the Weyl root system for each d-fold affinization matrix in terms of the root system of the corresponding Cartan matrix and some special null roots.

1. Introduction

Generalized intersection matrices were introduced by P. Slodowy in his study on isolated singularities (see [Slo1] or [Slo2]). An integral matrix $A = (A_{ij})_{i,j\in I}$ with the finite index set I is called a generalized intersection matrix or a GIM if for any $i, j \in I$, we have

$$\begin{aligned} &A_{ii} = 2, \\ &A_{ij} < 0 \quad \text{iff} \quad A_{ji} < 0, \\ &A_{ij} > 0 \quad \text{iff} \quad A_{ji} > 0. \end{aligned}$$

In this case, if A is symmetrizable, then we call it a symmetrizable intersection matrix or an SIM; and if A is symmetric, then we call it an intersection matrix or an IM. Clearly the notion of generalized intersection matrices is also a generalization of the notion of generalized Cartan matrices.

For any generalized intersection matrix, one can associate it to a root basis, as similar to what one dose for a generalized Cartan matrix. Therefore a natural question is how to describe or classify such kind of root bases.

When studying the elliptic singularities, K. Saito in [Sai] introduced the extended affine root systems, which are associated to positive semi-definite bilinear forms such that the rank of the radical of each such form is two. By using the diagrams with dotted edges he obtained a classification for these extended affine root systems.

B. Allison, N. Azam, S. Berman, Y. Gao and A. Pianzola in [AABGP] considered the extended affine root systems in general form, namely these extended affine root systems associated to arbitrary positive semi-definite bilinear forms without any restriction on the rank of the radical of each such form. They used the lattices or semi-lattices to obtain a classification for these extended affine root systems.

In this paper we consider symmetrizable intersection matrices and their root systems. The main method is applying braid-transformations on root bases defined

---

2000 *Mathematics Subject Classification.* Primary 17B22; Secondary 17B67.

Supported partially by the National Natural Science Foundation of China (No. 10325107), and the 973 Program (No. 2006CB805905).

by Slodowy and diagrams with dotted edges to our study. We state our main results as follows.

Firstly, inspired by the Auslander-Platzeck-Reiten tilts or co-tilts in representation theory of algebras we define the APR-transformation on root bases as a composition of certain braid-transformations. Using APR-transformations we prove that any positive semi-definite symmetrizable intersection matrix A with $\operatorname{corank} A \leq 1$ is braid-equivalent to a Cartan matrix or an affine generalized Cartan matrix. In this case, as a consequence we show that the intersection matrix Lie algebra of A in sense of Slodowy is isomorphic to a semi-simple or affine Kac-Moody Lie algebra.

Secondly, given any Cartan Matrix, we define its (standard) d-fold affinization matrix by adding some single roots or double roots of this Cartan matrix. This generalizes the corresponding notion of S. Berman and R. V. Moody [BM] and that of G. Benkart and E. Zelmanov [BZ], where they only considered the case that single roots were added. Furthermore, we prove that any positive semi-definite symmetrizable intersection matrix is braid-equivalent to a d-fold affinization matrix. By introducing the type of a d-fold affinization matrix and using standard d-fold affinization matrices, we give a complete classification of all positive semi-definite symmetrizable intersection matrices, up to braid-equivalence.

Finally, we give an explicit structure of the Weyl root system for each d-fold affinization matrix in terms of the root system of the corresponding Cartan matrix and some special null roots.

We should mention that in this paper we do not deal with another question how to define a suitable Lie algebra associated to a generalized intersection matrix just like the Kac-Moody Lie algebra associated to a generalized Cartan matrix. Usually this is a hard question. There were many attempts to do it. For example, Slodowy in [Slo2] defined generalized intersection matrix Lie algebras and intersection matrix Lie algebra; Berman and Moody in [BM] and Benkart and Zelmanov in [BZ] studied and classified the Lie algebras graded by finite root systems; Allison, Azam, Berman, Gao and Pianzola in [AABGP] considered the constructions of Lie algebras to realize the extended affine root systems; Saito and Yoshii in [SY] defined simply-laced elliptic Lie algebras by finite many generators and finite many relations depended on the classification of elliptic root systems given by Saito in [Sai]; and so on. In [XP], we also made an attempt to define the SIM Lie algebras for symmetrizable intersection matrices.

2. SIM-Root Bases and APR-Transformations

2.1. **GIM-Root Bases, SIM-Root Bases and Reflection Transformations.** Let us first recall the definitions of generalized intersection matrices, intersection matrices, GIM-root bases and braid-equivalences given by P. Slodowy in [Slo1], [Slo2]. Then we consider symmetrizable generalized intersection matrices, which are our emphasis.

Definition 2.1. Let $A = (A_{ij})_{i,j\in I}$ be an integral matrix, where I is a finite set. If the integers A_{ij} satisfies the following properties: for all $i, j \in I$,

$$\begin{aligned} &A_{ii} = 2, \\ &A_{ij} < 0 \quad \Leftrightarrow \quad A_{ji} < 0, \\ &A_{ij} > 0 \quad \Leftrightarrow \quad A_{ji} > 0, \end{aligned}$$

then we call A a *generalized intersection matrix*, or GIM for short.

If A is symmetric, we call A an *intersection matrix*, or IM for short.

Note that any generalized Cartan matrix is a generalized intersection matrix.

Definition 2.2. Let $A = (A_{ij})_{i,j\in I}$ be a GIM. A *generalized intersection matrix root basis* of A, or a GIM-root basis for short, is a triple (H, ∇, Δ) consisting of
(1) a finite-dimensional $\mathbb{Q}$-vector space H;
(2) $\nabla = \{h_i | i \in I\} \subseteq H$;
(3) $\Delta = \{\alpha_i | i \in I\} \subseteq H^* = \mathrm{Hom}_{\mathbb{Q}}(H, \mathbb{Q})$, such that

$$\alpha_j(h_i) = A_{ij} \quad \text{for all} \;\; i, j \in I.$$

We also call the generalized intersection matrix $A = (A_{ij})_{i,j\in I}$ is *the structural matrix* of the GIM-root basis (H, ∇, Δ).

Let $A = (A_{ij})_{i,j\in I}$ be a GIM, and (H, ∇, Δ) be the corresponding GIM-root basis. The *reflection transformations* are defined as follows: for every $\alpha \in \Delta$,

$$s_\alpha(h) := h - \alpha(h)h_\alpha, \quad h \in H.$$

The contragredient action of s_α on H^* is given by

$$s_\alpha(\gamma) := \gamma - \gamma(h_\alpha)\alpha, \quad \gamma \in H^*.$$

Here if $\alpha = \alpha_i$, we denote $h_\alpha = h_i$.

Two GIM-root bases (H, ∇, Δ) and (H, ∇', Δ') are called *braid-equivalent* if they can be transformed into each other by a sequence of transformations of the form

$$(H, \nabla, \Delta) \mapsto \cdots \mapsto (H, \nabla_k, \Delta_k) \mapsto (H, \nabla_{k+1}, \Delta_{k+1}) \mapsto \cdots \mapsto (H, \nabla', \Delta'),$$

where

$$\begin{array}{rcl} \Delta_{k+1} & = & (\Delta_k \setminus \{\beta\}) \cup \{s_\alpha(\beta)\}, \\ & & \qquad\qquad\qquad\qquad \text{for some} \quad \alpha, \beta \in \Delta_k. \\ \nabla_{k+1} & = & (\nabla_k \setminus \{h_\beta\}) \cup \{s_\alpha(h_\beta)\}, \end{array}$$

It is easy to see that braid-equivalence is an equivalence relation.

In the following we consider the symmetrizable case.

Let $A = (A_{ij})_{i,j\in I}$ be a *symmetrizable generalized intersection matrix*, that is, A is a generalized intersection matrix and there exists an invertible diagonal matrix $D = \mathrm{diag}(d_i)_{i\in I}$, where all d_i's are positive integers and $\gcd(d_i)_{i\in I} = 1$, such that DA is a symmetric matrix. In this case D is called the *symmetrizer* of A. Denote $DA := B = (B_{ij})_{i,j\in I}$. Let H be a $\mathbb{Q}$-vector space, and $\Delta := \{\alpha_i | i \in I\} \subseteq H$ be a linearly independent subset. Then there exists a symmetric bilinear form

$$(-,\,-) : H \times H \to \mathbb{Q}$$

such that

$$(\alpha_i,\, \alpha_j) = B_{ij} \quad \text{for } i, j \in I.$$

It is easy to see that

$$A_{ij} = \frac{2(\alpha_i,\, \alpha_j)}{(\alpha_i,\, \alpha_i)} \quad \text{for } i, j \in I$$

and

$$d_i = (\alpha_i, \alpha_i)/2 \quad \text{for } i \in I.$$

We call (H, Δ) an *SIM-root basis* of the symmetrizable generalized intersection matrix A, and A is called the structural matrix of (H, Δ). Note that since we have the above symmetric bilinear form, we can take $\nabla = \{\frac{2\alpha_i}{(\alpha_i,\, \alpha_i)} | i \in I\}$. Therefore we have no need to distinguish H and its dual space H^*. In this case, we have the following reflection transformations: for every $\alpha_i \in \Delta$,

$$s_{\alpha_i}(h) := h - \frac{2(\alpha_i,\, h)}{(\alpha_i,\, \alpha_i)}\alpha_i, \qquad h \in H.$$

Particularly, for all $\alpha_i,\ \alpha_j \in \Delta$,

$$s_{\alpha_i}(\alpha_j) := \alpha_j - A_{ij}\alpha_i.$$

From now on we call a symmetrizable generalized intersection matrix just a *symmetrizable intersection matrix*, or SIM for short. We always suppose $A = (A_{ij})_{i,j\in I}$ is an SIM. Accordingly, let H, Δ and the symmetric bilinear form $(-,-)$ be as above. Moreover, we suppose H is a $\mathbb{Q}$-vector space of dimension $|I|$. So Δ is a basis of H. In this case we say Δ is an SIM-root basis of the symmetrizable intersection matrix A, and A is the structural matrix of Δ.

If $A = (A_{ij})_{i,j\in I}$ is a (symmetrizable) generalized Cartan matrix, we call the corresponding SIM-root basis or GIM-root basis a *GCM-root basis.*

We fix the following notations and notions.

Let $\mathcal{W} \subseteq GL(H)$ be the subgroup of $GL(H)$ generated by s_{α_i}, $\alpha_i \in \Delta$. We call $\mathcal{W}$ the *Weyl group* of the SIM-root basis Δ, and

$$R^W := \{s(\alpha) \,|\, s \in \mathcal{W}, \alpha \in \Delta\}$$

the *Weyl root system* of Δ. Set

$$\Gamma := \sum_{i\in I} \mathbf{Z}\alpha_i.$$

The lattice Γ is called the *root lattice* of Δ.

If an element $\alpha \in H$ is non-isotropic, that is $(\alpha,\, \alpha) \neq 0$, then we define the dual $\alpha^\vee$ of α and the reflection $s_\alpha \in GL(H)$ as follows:

$$\alpha^\vee := \frac{2}{(\alpha,\, \alpha)}\alpha,$$

$$s_\alpha(\beta) := \beta - (\beta,\, \alpha^\vee)\alpha, \quad \text{for all } \beta \in H.$$

It is easy to check that

$$\alpha^{\vee\vee} = \alpha,$$

$$s_\alpha = s_{\alpha^\vee}, \quad s_\alpha^2 = id_H,$$

$$(s_\alpha(\beta),\, s_\alpha(\gamma)) = (\beta,\, \gamma), \quad s_\alpha(\beta)^\vee = s_\alpha(\beta^\vee).$$

2.2. **Braid-transformations and APR-transformations.** Let Δ be an SIM-root basis with a symmetrizable intersection matrix $A = (A_{ij})_{i,j\in I}$ as its structural matrix.

For all $a, b \in I$, we define an 'operator' $\tau_{a,b}$ on the root basis Δ as follows:

$$\tau_{a,b}\Delta = \Delta'$$

where $\Delta' = \{\, \alpha'_i \,|\, i \in I\}$ is defined as:

$$\begin{aligned} &\alpha'_i = \alpha_i && \text{for all } b \neq i \in I, \\ &\alpha'_b = s_{\alpha_a}(\alpha_b) = \alpha_b - A_{ab}\alpha_a. && \end{aligned}$$

The structural matrix of Δ' is $A' = (A'_{ij})_{i,j\in I}$, where

$$\begin{array}{ll} A'_{ij} = A_{ij} & \text{for } \ i \neq b, j \neq b, \\ A'_{ib} = A_{ib} - A_{ia}A_{ab} & \text{for } \ i \neq b, \\ A'_{bj} = A_{bj} - A_{ba}A_{aj} & \text{for } \ j \neq b, \\ A'_{bb} = 2. & \end{array}$$

It is easy to see that A' is also a symmetrizable intersection matrix and its symmetrizer is the same as that of A. Therefore, Δ' is also an SIM-root basis. We call $\tau_{a,b}$ a *braid-transformation*, and write $\tau_{a,b}A := A'$.

Two SIM-root bases Δ and Δ' are called *braid-equivalent*, if there exist $a_1, b_1, \cdots, a_t, b_t \in I$ such that

$$\tau_{a_t,b_t} \cdots \tau_{a_1,b_1} \Delta = \Delta'.$$

Obviously, the braid-transformation $\tau_{a,b}$ is invertible, and its inverse is also $\tau_{a,b}$. Therefore the braid-equivalence relation is an equivalence relation. In fact, the definition of the braid-equivalences of SIM-root bases given here is coincident with that of the corresponding braid-equivalences of GIM-root bases.

If the structural matrices of the above two SIM-root bases are respectively $A = (A_{ij})_{i,j\in I}$ and $A' = (A'_{ij})_{i,j\in I}$, we also say A and A' braid-equivalent. It is easy to show that a braid-transformation preserves the property of being symmetric, symmetrizable or indecomposable as well as the rank of a matrix.

Remark 2.3. In our definition, the structural matrix $A = (A_{ij})_{i,j\in I}$ of an SIM-root basis or a GIM-root basis is independent of the order of the finite index set I. Therefore two symmetrizable intersection matrices $A = (A_{ij})_{i,j\in I}$ and $A' = (A'_{ij})_{i,j\in I}$ of the same order of the finite index set are braid-equivalent if and only if A can be transformed into A' by a sequence of 'operators' of this form $\tau_{a,b}(a, b \in I)$ and permutations of elements in the index set I.

More generally, for $a \in I$ and a nonempty subset N of I, we define the operator $\tau_{a,\,N}$ as follows:

$$\tau_{a,\,N}\Delta = \Delta',$$

where $\Delta' = \{\alpha'_i \mid i \in I\}$ with

$$\begin{array}{ll} \alpha'_i = \alpha_i, & \text{for all } \ i \in I \setminus N; \\ \alpha'_i = s_{\alpha_a}(\alpha_i), & \text{for all } \ i \in N. \end{array}$$

The operator $\tau_{a,\,N}$ has the following property:

Proposition 2.4. *Let* $N = \{b_1, b_2, \cdots, b_t\} \subseteq I$. *Then*

$$\tau_{a,\,N}\Delta = \tau_{a,\,b_t} \cdots \tau_{a,\,b_1}\Delta.$$

Proof. We prove it by induction. For $t = 1$, it is clear. For $t > 1$, put $N' = N \setminus \{b_t\}$ and suppose that $\tau_{a,\,N'}\Delta = \tau_{a,\,b_{t-1}} \cdots \tau_{a,\,b_1}\Delta$. Set $\tau_{a,\,N'}\Delta = \Delta'$. Then $\alpha'_i = \alpha_i$ if $i \in I\backslash N'$, and $\alpha'_i = s_{\alpha_a}(\alpha_i)$ if $i \in N'$. Set $\tau_{a,\,b_t}\Delta' = \Delta''$. Then $\alpha''_i = \alpha'_i$ if $i \neq b_t$, and $\alpha''_{b_t} = s_{\alpha'_a}(\alpha'_{b_t}) = \alpha'_{b_t} - (\alpha'^{\vee}_a, \alpha'_{b_t})\alpha'_a$. Thus if $i \in I\backslash N$, then $\alpha''_i = \alpha_i$. If $i \in N'$, then $\alpha''_i = s_{\alpha_a}(\alpha_i)$. So, when $a \in I \setminus N'$, we have $\alpha''_{b_t} = \alpha_{b_t} - (\alpha^{\vee}_a, \alpha_{b_t})\alpha_a = s_{\alpha_a}(\alpha_{b_t})$. And when $a \in N'$, we have $\alpha''_{b_t} = \alpha_{b_t} - (-\alpha^{\vee}_a, \alpha_{b_t})(-\alpha_a) = s_{\alpha_a}(\alpha_{b_t})$. Therefore we have $\tau_{a,\,N}\Delta = \Delta'' = \tau_{a,\,b_t}\Delta' = \tau_{a,\,b_t} \cdots \tau_{a,\,b_1}\Delta$. □

Definition 2.5. Let $A = (A_{ij})_{i,j\in I}$ be an SIM, and Δ be the SIM-root basis of A. Given $a \in I$, let $N(a) = \{a\} \cup \{b \in I |\ A_{ab} < 0\}$. We call $\tau_a := \tau_{a,\,N(a)}$ an

Auslander-Platzeck-Reiten transformation of the root basis Δ at the point a, or an APR-transformation at a for brevity.

Remark 2.6. The notion of Auslander-Platzeck-Reiten transformations is inspired by the notion of APR-tilts and APR-co-tilts given by M. Auslander, M. I. Platzeck and I. Reiten in [APR], which plays an important role in the representation theory of algebras. From the view of tilting theory in the representation theory of algebras, L. Peng in [Peng] gave an explicit explanation of the relation between APR-transformations given here and APR-tilts as well as APR-co-tilts.

Proposition 2.4 says that an APR-transformation is a composite of certain braid-transformations. In the following we shall produce a formula to compute the structural matrix of an APR-transformation of an SIM-root basis. This formula is very useful for our computations.

Proposition 2.7. *Let τ_a be an APR-transformation. Denote $\Delta^a := \tau_a \Delta$ with structural matrix A^a $(:= \tau_a A)$, where $A^a = (A^a_{ij})_{i,j\in I}$ is the matrix such that for all $i, j \in I$, $A^a_{ij} = (\alpha^a_j,\ (\alpha^a_i)^\vee)$. Then we have*

$$\begin{array}{ll} A^a_{ij} = A_{ij} & \textit{for} \ \ i,j \notin N(a) \ \ \textit{or} \ \ i,j \in N(a), \\ A^a_{ij} = A_{ij} - A_{ia}A_{aj} & \textit{otherwise.} \end{array}$$

Proof. By definition, for $i \notin N(a)$, we have $\alpha^a_i = \alpha_i$; for $i \in N(a)$, we have $\alpha^a_i = s_{\alpha_a}(\alpha_i) = \alpha_i - A_{ai}\alpha_a$. So when $i, j \notin N(a)$ we have

$$A^a_{ij} = (\alpha^a_j,\ (\alpha^a_i)^\vee) = (\alpha_j,\ \alpha_i^\vee) = A_{ij}.$$

When $i, j \in N(a)$ we have

$$A^a_{ij} = (s_{\alpha_a}(\alpha_j),\ s_{\alpha_a}(\alpha_i)^\vee) = (\alpha_j,\ \alpha_i^\vee) = A_{ij}.$$

When $i \in N(a)$ and $j \notin N(a)$ we have

$$\begin{aligned} A^a_{ij} &= (\alpha_j,\ s_{\alpha_a}(\alpha_i)^\vee) = \frac{2(\alpha_j,\ \alpha_i - A_{ai}\alpha_a)}{(s_{\alpha_a}(\alpha_i),\ s_{\alpha_a}(\alpha_i))} \\ &= A_{ij} - \frac{2A_{ai}(\alpha_j,\ \alpha_a)}{(\alpha_i,\ \alpha_i)} = A_{ij} - \frac{4(\alpha_i,\ \alpha_a)(\alpha_j,\ \alpha_a)}{(\alpha_a,\ \alpha_a)(\alpha_i,\ \alpha_i)} = A_{ij} - A_{ia}A_{aj}. \end{aligned}$$

When $i \notin N(a)$ and $j \in N(a)$, we have

$$A^a_{ij} = (s_{\alpha_a}(\alpha_j),\ \alpha_i^\vee) = \frac{2(\alpha_j - A_{aj}\alpha_a,\ \alpha_i)}{(\alpha_i,\ \alpha_i)} = A_{ij} - \frac{2A_{aj}(\alpha_a,\ \alpha_i)}{(\alpha_i,\ \alpha_i)} = A_{ij} - A_{ia}A_{aj}.$$

□

By the definition of $N(a)$ and the above formula, we have the following immediate consequence.

Corollary 2.8. *If A is a symmetrizable generalized Cartan matrix, then for any APR-transformation τ_a, we have $\tau_a A = A$.* □

2.3. **SIM's and Dynkin Diagrams.**

For any symmetrizable intersection matrix $A = (A_{ij})_{i,j\in I}$ we define a diagram $D(A)$ associated to A, called the Dynkin diagram of A, as follows.

Let I be the vertex set. If $a_{ij} \neq 0$ and $a_{ij} \neq a_{ji}$, the vertices i and j are connected by a solid edge if $a_{ij} < 0$ or by a dotted edge if $a_{ij} > 0$, and these edges are equipped with an ordered pair of integers $(|a_{ij}|, |a_{ji}|)$. If $a_{ij} = a_{ji}$, the vertices

i and j are connected by $(-a_{ij})$ solid edges if $a_{ij} \leq 0$ or by a_{ij} dotted edges if $a_{ij} > 0$. Here, our definition is slightly different from that given by Slodowy in [Slo1] and [Slo2]. Our edges are valued, while those defined by Slodowy are not valued.

For two SIM's A and A', $D(A)$ and $D(A')$ are said to be braid-equivalent if their SIM-root bases are braid-equivalent. In addition, if $\tau_{a,\,b}A = A'$, then we also write $\tau_{a,\,b}D(A) = D(A')$.

A diagram without dotted edges is called a *solid diagram.* It is clear that $D(A)$ is a solid diagram if and only if A is a generalized Cartan matrix. In this case $D(A)$ is invariant under any APR-transformation.

For the sake of convenience, we label all Dynkin diagrams of positive-definite GCM's (that is, classic Dynkin diagrams) as follows:

A type:



B type:



C type:



D type:



E_6 type:



E_7 type:



E_8 type:



F_4 type:



G_2 type:



Let a_t be the left end point in each Dynkin diagram as above, for example, $a_t = a_2$ in E_6 case, similarly for others. Then we have

Lemma 2.9. *Let $A = (A_{ij})_{i,j\in I}$ be an SIM such that A is positive semi-definite and* $\operatorname{corank} A \le 1$. *If there is a $d \in I$ such that the full subdiagram $D(A) \setminus \{d\}$ of $D(A)$ is a Dynkin diagram of type $A_l, B_l, C_l, D_l, E_6, E_7, E_8, F_4, G_2$ with $A_{di} \le 0$ for all $i \in I \setminus \{d, a_t\}$, then the SIM-root basis Δ is braid-equivalent to a GCM-root basis by successive APR-transformations at some vertices in $I \setminus \{d\}$.*

Proof. If $A_{da_t} \le 0$, then A itself is a GCM. So we can assume that $A_{da_t} > 0$. Note that A is positive semi-definite. Thus we have $A_{da_t}A_{a_td} \le 4$ and $h \le 4$, where $h = |\sum\limits_{i\in I\setminus\{d,a_t\}} A_{di}|$.

In the following diagrams, the valuation on the vertical edge connecting the top vertex i and the bottom vertex j is denoted (a_{ij}, a_{ji}). In other cases, (a_{ij}, a_{ji}) means the valuation on the edge linking the left vertex i and the right vertex j.

Since $A_{da_t}A_{a_td} \le 4$, we have $A_{da_t} \le 4$. So we verify the statement case by case, according to the value of A_{da_t} being equal to $4, 3, 2$ or 1, respectively. We will not list some obviously impossible cases, taking into account the condition that our matrix is symmetrizable and positive semi-definite.

Case 1. Assume that $A_{da_t} = 4$. Then we have $A_{a_td} = 1$. There are three sub-cases.

(1.1) Let $D(A) \setminus \{d\}$ be of type A_l.

It is clear that $D(A) \setminus \{d\}$ must be of type A_1. Thus $\tau_{a_1}\Delta$ is a GCM-root basis of type $A_2^{(2)}$.

(1.2) Assume that $D(A) \setminus \{d\}$ is of type B_l.

It is easy to see there is only one case:



The diagram of $\tau_b\tau_{a_0}\Delta$ is



(1.3) $D(A) \setminus \{d\}$ can not be of type $C_l (l > 2), D_l, E_6, E_7, E_8, F_4, G_2$.

Case 2. Assume that $A_{da_t} = 3$. Then we have $A_{a_td} = 1$. There are also three sub-cases.

(2.1) Let $D(A) \setminus \{d\}$ be of type A_l.

It is easy to see that $D(A) \setminus \{d\}$ must be of type A_1 or A_2.

If it is A_1, $\tau_{a_1}\Delta$ is a GCM-root basis of type G_2.

If it is A_2, there are only two possibilities. One is



The diagram of $\tau_{a_1}\tau_{a_2}\Delta$ is



Another one is



The diagram of $\tau_{a_2}\Delta$ is



(2.2) $D(A)\setminus\{d\}$ can not be of type $B_l, C_l, D_l, E_6, E_7, E_8, F_4$.

(2.3) Suppose $D(A)\setminus\{d\}$ is of type G_2. There are only two possibilities. The first one is



The diagram of $\tau_b\tau_{a_0}\tau_b\tau_{a_0}\Delta$ is



The second one is



The diagram of $\tau_b\tau_{a_0}\Delta$ is



Case 3. Assume that $A_{da_t} = 2$ and $A_{a_t d} = 1$. Then we have the following six sub-cases.

(3.1) Suppose $D(A)\setminus\{d\}$ is of type A_l. There are two possibilities, according to the value of h.

(3.1.1) If $h = 0$, the diagram of Δ is



The diagram of $\tau_{a_1}\tau_{a_2}\cdots\tau_{a_t}\Delta$ is



(3.1.2) If $h = 2$, the diagram of Δ is



The diagram of $\tau_{a_{i+1}}\tau_{a_{i+2}}\cdots\tau_{a_t}\Delta$ is



(3.2) Assume that $D(A) \setminus \{d\}$ is of type B_l. Then there are three possibilities.

(3.2.1) If $h = 0$, the diagram of Δ is



The diagram of $\tau_{a_t}\cdots\tau_{a_0}\tau_b\tau_{a_0}\cdots\tau_{a_t}\Delta$ is



(3.2.2) If $h = 1$, the diagram of Δ is only one possibility:



The diagram of $\tau_b\tau_{a_0}\cdots\tau_{a_t}\Delta$ is



(3.2.3) If $h = 2$, then there are two possibilities. One is



The diagram of $\tau_{a_t}\Delta$ is



Another one is



The diagram of $\tau_{a_0}\Delta$ is



(3.3) Suppose $D(A)\setminus\{d\}$ is of type C_l. Similarly, according to the value of h, we have the following cases.

(3.3.1) If $h=0$, the diagram of Δ is



The diagram of $\tau_{a_t}\cdots\tau_{a_0}\tau_b\tau_{a_0}\cdots\tau_{a_t}\Delta$ is



(3.3.2) If $h=2$, the diagram of Δ is only one possibility:



The diagram of $\tau_{a_t}\Delta$ is



(3.3.3) If $h=4$, the diagram of Δ is only one possibility:



The diagram of $\tau_{a_0}\Delta$ is



(3.4) Assume that $D(A)\setminus\{d\}$ is of type D_l. We have the following cases.

(3.4.1) If $h=0$, the diagram of Δ is



The diagram of $\tau_{a_t}\cdots\tau_{a_0}\tau_c\tau_b\tau_{a_0}\cdots\tau_{a_t}\Delta$ is



(3.4.2) If $h = 2$, there are two cases. One is



The diagram of $\tau_{a_t}\Delta$ is



Another one is



The diagram of $\tau_b\tau_{a_0}\tau_{a_1}\Delta$ is



(3.5) $D(A) \setminus \{d\}$ can not be of type $E_l (l = 6, 7, 8)$ or G_2.

(3.6) Suppose $D(A) \setminus \{d\}$ is of type F_4. Then h must be equal to 1 and there are only two possibilities. One is



The diagram of $\tau_{b_2}\tau_{b_1}\tau_{a_0}\tau_{a_1}\Delta$ is



Another one is



The diagram of $\tau_{b_2}\tau_{b_1}\tau_{a_0}\tau_{a_1}\Delta$ is



Case 4. Assume that $A_{da_t} = 2$ and $A_{a_td} = 2$. Then we have the following sub-cases.

(4.1) Assume that $D(A) \setminus \{d\}$ is of type A_l.

If it is A_1, $\tau_{a_1}\Delta$ is a GCM-root basis of type $A_1^{(1)}$.

If $l > 1$, then h must be equal to 1. There is only one possibility



The diagram of $\tau_{a_1} \cdots \tau_{a_t}\Delta$ is



(4.2) Assume that $D(A)\setminus\{d\}$ is of type B_l. Then h must be equal to 1 and there are two possibilities. One is



The diagram of $\tau_{a_t} \cdots \tau_{a_0}\tau_b\tau_{a_0} \cdots \tau_{a_t}\Delta$ is



Another one is



The diagram of $\tau_b\tau_{a_0}\Delta$ is



(4.3) Suppose $D(A) \setminus \{d\}$ is of type C_l.

(4.3.1) If $h = 1$, the diagram of Δ is only one possibility:



The diagram of $\tau_{a_t}\cdots\tau_{a_0}\tau_b\tau_{a_0}\cdots\tau_{a_t}\Delta$ is



(4.3.2) If $h = 2$, the diagram of Δ is only one possibility:



The diagram of $\tau_b\tau_{a_0}\Delta$ is



(4.4) Assume that $D(A)\setminus\{d\}$ is of type D_l. Then the diagram of Δ is only one possibility:



The diagram of $\tau_{a_{t-1}}\cdots\tau_{a_0}\tau_c\tau_b\tau_{a_0}\cdots\tau_{a_t}\Delta$ is



(4.5) Suppose $D(A)\setminus\{d\}$ is of type E_l. The diagram of Δ is only one possibility:



where $t = 2, 3, 4$.

For $t = 2$, the diagram of $\tau_c\tau_{a_0}\tau_{b_1}\tau_{a_1}\tau_{a_0}\tau_c\tau_{b_2}\tau_{b_1}\tau_{a_0}\tau_{a_1}\cdots\tau_{a_t}\Delta$ is



For $t = 3$, the diagram of $\tau_{b_2}\tau_{b_1}\tau_{a_0}\tau_{a_1}\tau_c\tau_{a_0}\tau_{a_2}\tau_{a_1}\tau_{b_1}\tau_{a_0}\tau_c\tau_{b_2}\tau_{b_1}\tau_{a_0}\tau_{a_1}\cdots\tau_{a_t}\Delta$ is



For $t = 4$, the diagram of $\tau_{a_4}\tau_{a_3}\tau_{a_2}\tau_{a_1}\tau_{a_0}\tau_{b_1}\tau_c\tau_{a_0}\tau_{a_1}\tau_{b_2}\tau_{b_1}\tau_{a_0}\tau_{a_2}\tau_{a_1}\tau_c\tau_{a_0}\tau_{a_3}\tau_{a_2}\tau_{a_1}$ $\tau_{b_1}\tau_{a_0}\tau_c\tau_{b_2}\tau_{b_1}\tau_{a_0}\tau_{a_1}\cdots\tau_{a_t}\Delta$ is



(4.6) Assume that $D(A) \setminus \{d\}$ is of type F_4. The diagram of Δ is only one possibility:



The diagram of $\tau_{a_1}\tau_{a_0}\tau_{b_1}\tau_{b_2}\tau_{a_0}\tau_{b_1}\tau_{a_0}\tau_{a_1}\Delta$ is



(4.7) Assume that $D(A) \setminus \{d\}$ is of type G_2. And the diagram of Δ is only one possibility:



The diagram of $\tau_{a_0}\tau_b\tau_{a_0}\Delta$ is



Case 5. Assume that $A_{da_t} = 1$ and $A_{a_td} = 1$. Then we have the following sub-cases.

(5.1) Assume that $D(A) \setminus \{d\}$ is of type A_l. Then we have the following three cases.

(5.1.1) If $h = 0$, the diagram of Δ is



The diagram of $\tau_{a_1}\cdots\tau_{a_{t-1}}\tau_{a_t}\Delta$ is



(5.1.2) If $h = 1$, the diagram of Δ is



The diagram of $\tau_{a_{i+1}}\tau_{a_{i+2}}\cdots\tau_{a_t}\Delta$ is



(5.1.3) If $h = 2$, there are two cases. One is



The diagram of $\tau_{a_t}\Delta$ is



Another one is



The diagram of $\tau_{a_2}\Delta$ is



(5.2) Assume that $D(A)\setminus\{d\}$ is of type B_l. Then similarly we have the following cases.

(5.2.1) If $h = 0$, the diagram of Δ is



The diagram of $\tau_{a_t}\cdots\tau_{a_0}\tau_b\tau_{a_0}\cdots\tau_{a_t}\Delta$ is



(5.2.2) If $h = 1$, then there are two possibilities. One is



The diagram of $\tau_{a_{i+1}}\tau_{a_{i+2}}\cdots\tau_{a_t}\Delta$ is



Another one is



The diagram of $\tau_{a_0}\cdots\tau_{a_t}\Delta$ is



(5.3) Assume that $D(A)\setminus\{d\}$ is of type C_l. Then we have the following cases.

(5.3.1) If $h = 0$, the diagram of Δ is



The diagram of $\tau_{a_t}\cdots\tau_{a_0}\tau_b\tau_{a_0}\cdots\tau_{a_t}\Delta$ is



(5.3.2) If $h = 1$, the diagram of Δ is only one possibility:



The diagram of $\tau_{a_{i+1}}\tau_{a_{i+2}}\cdots\tau_{a_t}\Delta$ is



(5.3.3) If $h = 2$, the diagram of Δ is only one possibility:



The diagram of $\tau_{a_0} \cdots \tau_{a_t} \Delta$ is



(5.4) Suppose $D(A) \setminus \{d\}$ is of type D_l. As before we have the following cases according to the value of h.

(5.4.1) If $h = 0$, the diagram of Δ is



The diagram of $\tau_{a_t} \cdots \tau_{a_1} \tau_{a_0} \tau_b \tau_c \tau_{a_0} \tau_{a_1} \cdots \tau_{a_t} \Delta$ is



(5.4.2) If $h = 1$, then there are two possibilities. One is



The diagram of $\tau_{a_{i+1}} \tau_{a_{i+2}} \cdots \tau_{a_t} \Delta$ is



Another one is



The diagram of $\tau_b\tau_{a_0}\tau_{a_1}\tau_{a_2}\cdots\tau_{a_t}\Delta$ is



(5.5) Suppose $D(A)\setminus\{d\}$ is of type E_l. We have the following cases.

(5.5.1) If $h=0$, the diagram of Δ is



where $t=2,3,4.$

For $t=2$, the diagram of $\tau_{b_2}\tau_{b_1}\tau_{a_0}\tau_{a_1}\tau_c\tau_{a_0}\tau_{a_t}\cdots\tau_{a_1}\tau_{b_1}\tau_{a_0}\tau_{b_2}\tau_{b_1}\tau_c\tau_{a_0}\tau_{a_1}\cdots\tau_{a_t}\Delta$ is



For $t=3$, the diagram of $\tau_{a_3}\tau_{a_2}\tau_{a_1}\tau_{a_0}\tau_{b_1}\tau_c\tau_{a_0}\tau_{a_1}\tau_{b_2}\tau_{b_1}\tau_{a_0}\tau_{a_2}\tau_{a_1}\tau_c\tau_{a_0}\tau_{a_t}\cdots\tau_{a_1}\tau_{b_1}$ $\tau_{a_0}\tau_{b_2}\tau_{b_1}\tau_c\tau_{a_0}\tau_{a_1}\cdots\tau_{a_t}\Delta$ is



For $t=4$, the diagram of $\tau_{a_4}\tau_{a_3}\tau_{a_2}\tau_{a_1}\tau_{a_0}\tau_{b_1}\tau_c\tau_{a_0}\tau_{a_1}\tau_{b_2}\tau_{b_1}\tau_{a_0}\tau_{a_2}\tau_{a_1}\tau_c\tau_{a_0}\tau_{a_3}\tau_{a_2}\tau_{a_1}$ $\tau_{b_1}\tau_{a_0}\tau_{a_4}\tau_{a_3}\tau_{a_2}\tau_{a_1}\tau_c\tau_{b_2}\tau_{b_1}\tau_{a_0}\tau_{a_1}\tau_{b_1}\tau_c\tau_{a_0}\tau_{a_2}\tau_{a_1}\tau_{b_2}\tau_{b_1}\tau_{a_0}\tau_{a_3}\tau_{a_2}\tau_{a_1}\tau_c\tau_{a_0}\tau_{a_t}\cdots\tau_{a_1}\tau_{b_1}\tau_{a_0}$ $\tau_{b_2}\tau_{b_1}\tau_c\tau_{a_0}\tau_{a_1}\cdots\tau_{a_t}\Delta$ is



(5.5.2) If $h=1$, then we have four cases. The first one is



where $t=2,3,4.$

The diagram of $\tau_{a_t}\Delta$ is



The second one is



where $t = 2, 3$.

The diagram of $\tau_{b_2}\tau_{b_1}\tau_{a_0}\tau_{a_1}\tau_{a_2}\cdots\tau_{a_t}\Delta$ is



The third one is



The diagram of $\tau_c\tau_{a_0}\tau_{a_1}\tau_{a_2}\Delta$ is



The last one is



where $t = 2, 3, 4$.

The diagram of $\tau_{a_t}\cdots\tau_{a_1}\tau_{a_0}\tau_{b_1}\tau_c\tau_{a_0}\tau_{a_1}\tau_{a_2}\cdots\tau_{a_t}\Delta$ is



(5.6) Suppose $D(A) \setminus \{d\}$ is of type F_4. Also we have the following cases.

(5.6.1) If $h = 0$, the diagram of Δ is



The diagram of $\tau_{a_1}\tau_{a_0}\tau_{b_1}\tau_{b_2}\tau_{a_0}\tau_{b_1}\tau_{a_1}\tau_{a_0}\tau_{b_1}\tau_{b_2}\tau_{a_1}\tau_{a_0}\tau_{b_1}\tau_{a_0}\tau_{a_1}\Delta$ is



(5.6.2) If $h = 1$, then there are two cases. One is



The diagram of $\tau_{a_1}\Delta$ is



Another one is



The diagram of $\tau_{a_1}\tau_{a_0}\tau_{b_1}\tau_{a_0}\tau_{a_1}\Delta$ is



(5.7) Finally, we assume that $D(A) \setminus \{d\}$ is of type G_2.

(5.7.1) If $h = 0$, the diagram of Δ is



The diagram of $\tau_{a_0}\tau_b\tau_{a_0}\tau_b\tau_{a_0}\Delta$ is



(5.7.2) If $h = 1$, the diagram of Δ is only one possibility:



The diagram of $\tau_{a_0}\Delta$ is



Case 6. Assume that $A_{da_t} = 1$ and $A_{a_t d} = 2$.
(6.1) Suppose $D(A) \setminus \{d\}$ is of type A_l.
(6.1.1) When $h = 0$, the diagram of Δ is



The diagram of $\tau_{a_1}\tau_{a_2}\cdots\tau_{a_t}\Delta$ is



(6.1.2) When $h = 1$, there are two cases. One is



The diagram of $\tau_{a_t}\Delta$ is



Another one is



The diagram of $\tau_{a_2}\tau_{a_3}\Delta$ is



(6.2) Suppose $D(A) \setminus \{d\}$ is of type B_l.
(6.2.1) When $h = 0$, the diagram of Δ is



The diagram of $\tau_{a_t}\cdots\tau_{a_0}\tau_b\tau_{a_0}\cdots\tau_{a_t}\Delta$ is



(6.2.2) When $h = 1$, there are two cases. One is



The diagram of $\tau_{a_t}\Delta$ is



Another one is



The diagram of $\tau_{a_0}\Delta$ is



(6.3) Suppose $D(A) \setminus \{d\}$ is of type C_l.

(6.3.1) When $h = 0$, the diagram of Δ is



The diagram of $\tau_{a_t} \cdots \tau_{a_0}\tau_b\tau_{a_0} \cdots \tau_{a_t}\Delta$ is



(6.3.2) When $h = 1$, there are two cases. One is



The diagram of $\tau_{a_t}\Delta$ is



Another one is



The diagram of $\tau_b\tau_{a_0} \cdots \tau_{a_t}\Delta$ is



(6.3.3) When $h = 2$, the diagram of Δ is only one possibility:



The diagram of $\tau_{a_0}\Delta$ is



(6.4) Suppose $D(A) \setminus \{d\}$ is of type D_l.
(6.4.1) When $h = 0$, the diagram of Δ is



The diagram of $\tau_{a_t} \cdots \tau_{a_0}\tau_c\tau_b\tau_{a_0} \cdots \tau_{a_t}\Delta$ is



(6.4.2) When $h = 1$, there are two cases. One is



The diagram of $\tau_{a_t}\Delta$ is



Another one is



The diagram of $\tau_b\tau_{a_0}\tau_{a_1}\Delta$ is



(6.5) Finally, $D(A) \setminus \{d\}$ can not be of type $E_l, F_4,$ or G_2.

Case 7. Assume that $A_{da_t} = 1$ and $A_{a_t d} = 3$.

(7.1) Suppose $D(A) \setminus \{d\}$ is of type A_l.

It is easy to see that $D(A) \setminus \{d\}$ must be of type A_1 or A_2.

If it is A_1, $\tau_{a_1}\Delta$ is a GCM-root basis of type G_2.

If it is A_2, there are only two cases. One is



The diagram of $\tau_{a_1}\tau_{a_2}\Delta$ is



Another one is



The diagram of $\tau_{a_2}\Delta$ is



(7.2) $D(A) \setminus \{d\}$ can not be of type $B_l, C_l, D_l, E_6, E_7, E_8, F_4,$ or G_2.

Case 8. Assume that $A_{da_t} = 1$ and $A_{a_t d} = 4$.

(8.1) Suppose $D(A) \setminus \{d\}$ is of type A_l.

It is clear that $D(A) \setminus \{d\}$ must be of type A_1. Thus $\tau_{a_1}\Delta$ is a GCM-root basis of type $A_2^{(2)}$.

(8.2) Suppose $D(A) \setminus \{d\}$ is C_l. We can easily see that there is only one case:



The diagram of $\tau_b\tau_{a_0}\Delta$ is



(8.3) $D(A) \setminus \{d\}$ can not be of type $B_l (l > 2), D_l, E_6, E_7, E_8, F_4,$ or G_2.

Up to now, we have discussed all possible cases and the proof is finished. □

Now we can show one of our main results.

Theorem 2.10. *Let $A = (A_{ij})_{i,j \in I}$ be an SIM such that A is positive semi-definite and* $\operatorname{corank} A \leq 1$. *Then for any $d \in I$ such that $(A_{ij})_{i,j \in I \setminus \{d\}}$ is positive definite, one can use APR-transformations at some vertices in $I \setminus \{d\}$ such that A is braid-equivalent to a GCM.*

Proof. By induction on $|I|$, we shall prove that for any $d \in I$, if $(A_{ij})_{i,j\in I\setminus\{d\}}$ is positive definite, then $D(A)$ is braid-equivalent to a solid diagram by successive APR-transformations at some vertices in $I \setminus \{d\}$.

Obviously $|I| > 1$, and there exists $d \in I$ such that $(A_{ij})_{i,j\in I\setminus\{d\}}$ is positive definite. Then by the induction hypothesis, the SIM-root basis Δ of A is braid-equivalent to an SIM-root basis Δ' with structural matrix $A' = (A'_{ij})_{i,j\in I}$ such that $D(A') \setminus \{d\}$ is a disjoint union of some classic Dynkin diagrams by successive APR-transformations at some vertices in $I \setminus \{d\}$.

If $D(A') \setminus \{d\}$ is not connected, then there exist two nonempty and disjoint subsets I_1 and I_2 of $I \setminus \{d\}$ such that $I \setminus \{d\} = I_1 \dot{\cup} I_2$ and $D(A') \setminus \{d\} = D(A'_1) \dot{\cup} D(A'_2)$, where $A'_t = (A'_{ij})_{i,j\in I_t}$, $t = 1, 2$. Thus there exist $a_1, \cdots, a_m \in I_1$ and $b_1, \cdots, b_n \in I_2$ such that both $\tau_{a_m} \cdots \tau_{a_1}(D(A'), I_1 \cup \{d\})$ and $\tau_{b_n} \cdots \tau_{b_1}(D(A'), I_2 \cup \{d\})$ are solid diagrams, where $(D(A'), I_t \cup \{d\})$ is the full subdiagram of $D(A')$ with vertex sets $I_t \cup \{d\}$, for $t = 1, 2$. Note that $D(A') \setminus \{d\}$ is a disjoint union of some classic Dynkin diagrams and so invariant under any APR-transformation at $i \in I \setminus \{d\}$. Thus $\tau_{b_n} \cdots \tau_{b_1} \tau_{a_m} \cdots \tau_{a_1}(D(A'))$ is a solid diagram. Therefore, $D(A)$ is braid-equivalent to a solid diagram by successive APR-transformations at some vertices in $I \setminus \{d\}$.

If $D(A') \setminus \{d\}$ is connected, then $D(A') \setminus \{d\}$ is the Dynkin diagram of type $A_l, B_l, C_l, D_l, E_6, E_7, E_8, F_4, G_2$. Then by the induction hypothesis, $D(A')$ is braid-equivalent to $D(A'')$ by the APR-transformations at some vertices in $I \setminus \{d, a_t\}$, where $A'' = (A''_{ij})_{i,j\in I}$ is an SIM such that $D(A'') \setminus \{a_t\}$ is a solid diagram. It is clear that $D(A'') \setminus \{d\} = D(A') \setminus \{d\}$. Therefore, by Lemma 2.9, $D(A'')$ is braid-equivalent to a solid diagram by successive APR-transformations at some vertices in $I \setminus \{d\}$. This implies that $D(A)$ is also braid-equivalent to a solid diagram by successive APR-transformations at some vertices in $I \setminus \{d\}$. □

Let us recall the notion and some results on IM-Lie algebras of Slodowy in [Slo1] and [Slo2].

Let $A = (A_{ij})_{i,j\in I}$ be an IM, Δ be the IM-root basis with structure matrix A, and $(-, -)$ be the symmetric bilinear form induced by A.

We first introduce an 'auxiliary Lie algebra' $\widetilde{\mathfrak{g}}(A)$, whose generators are $e_\alpha, \alpha \in \pm\Delta$ and $\mathfrak{h} = H \otimes_{\mathbb{Q}} \mathbb{C}$, and whose defining relations are as follows:

(A1) $[h, h'] = 0$, if $h, h' \in \mathfrak{h}$;

(A2) $[h, e_\alpha] = (h, \alpha)e_\alpha$, if $h \in \mathfrak{h}, \alpha \in \pm\Delta$;

(A3) $[e_\alpha, e_{-\alpha}] = -\alpha^\vee$, if $\alpha \in \pm\Delta$.

Thus, by defining $\deg e_\alpha = \alpha$ and $\deg h = 0$, the root lattice $\Gamma = \mathbf{Z} \cdot \Delta$ produces a gradation: $\widetilde{\mathfrak{g}}(A) = \bigoplus\limits_{\gamma\in\Gamma} \widetilde{\mathfrak{g}}_\gamma$.

Definition 2.11. (See [Slo1], [Slo2]) Let A be an intersection matrix, and $\mathfrak{r}$ be the ideal of $\widetilde{\mathfrak{g}}(A)$ generated by the following subspaces:

(IM4) $\widetilde{\mathfrak{g}}_\gamma$, where $\gamma \in \Gamma$ and $(\gamma, \gamma) > 2$.

We call the quotient Lie algebra $\mathrm{IM}(A) = \widetilde{\mathfrak{g}}(A)/\mathfrak{r}$ the *intersection matrix Lie algebra* associated to A, or IM-Lie algebra for brevity.

The following two results belong to Slodowy.

Proposition 2.12. *Let Δ and Δ' be two IM-bases with structure matrices A and A' respectively. If they are braid-equivalent, then IM(A) $\simeq$IM(A').* □

Proposition 2.13. *Let A be a symmetric generalized Cartan matrix. Then IM(A) is isomorphic to the Kac-Moody Lie algebra $\mathfrak{g}(A)$.* $\square$

By Proposition 2.12, Proposition 2.13 and Theorem 2.10, we have the following consequence.

Corollary 2.14. *Let A be a positive semi-definite IM with* $\operatorname{corank} A \leq 1$. *Then IM$(A)$ is isomorphic to a Kac-Moody Lie algebra.* $\square$

3. d-fold Affinizations and Classification of Positive Semi-Definite SIM's

In this section, we classify all positive semi-definite SIM's up to braid-equivalence. For this aim, we first introduce the notion of d-fold affinizations.

3.1. d-fold Affinizations.

Let C be an $l \times l$ Cartan matrix, $\dot{R}$ be the root system of C, and $\dot{\Delta} = \{\alpha_1, \alpha_2, \cdots, \alpha_l\} \subseteq \dot{R}$ be the root basis. Let $(-, -)_C$ be the symmetric bilinear form induced by C (so we have $C_{ij} = \frac{2(\alpha_i,\, \alpha_j)_C}{(\alpha_i,\, \alpha_i)_C}$). We define the d-fold affinizations of C as follows:

Definition 3.1. *A d-fold affinization of C* is an $(l + d)$-matrix $C^{[d]}$ of the form

$$C^{[d]}_{ij} = \frac{2(\alpha_i,\ \alpha_j)_C}{(\alpha_i,\ \alpha_i)_C}, \quad \text{where} \quad \alpha_{l+1}, \cdots, \alpha_{l+d} \in \dot{R} \cup 2\dot{R},$$

such that all $C^{[d]}_{ij}$ are integers. Here $2\dot{R} = \{2\alpha \mid \alpha \in \dot{R}\}$.

Particularly, if the above $\alpha_{l+1}, \cdots, \alpha_{l+d} \in \dot{\Delta} \cup 2\dot{\Delta}$, then we call the corresponding d-fold affinization matrix $C^{[d]}$ *standard*.

Remark 3.2. Our definition of d-fold affinization matrices is a generalization of that given by S. Berman and R. V. Moody in [BM] and by G. Benkart and E. Zelmanov in [BZ]. They only considered the case that $\alpha_{l+1}, \cdots, \alpha_{l+d} \in \dot{R}$.

Proposition 3.3. *Let C be a Cartan matrix and $C^{[d]}$ is a d-fold affinization of C. Then $C^{[d]}$ is a symmetrizable positive semi-definite matrix with* $\operatorname{corank} C^{[d]} = d$.

Proof. It is a direct checking. $\square$

In Definition 3.1, for the d-fold affinization matrix $C^{[d]}$, we have the SIM-root basis $\Delta = \{\alpha_1, \alpha_2, \cdots, \alpha_l, \beta_{l+1}, \cdots, \beta_{l+d}\}$, where $\beta_{l+1}, \cdots, \beta_{l+d}$ correspond respectively to $\alpha_{l+1}, \cdots, \alpha_{l+d} \in \dot{R} \cup 2\dot{R}$, that is $(\beta_{l+i}, \alpha_j)_{C^{[d]}} = (\alpha_{l+i}, \alpha_j)_C$, $(\beta_{l+i}, \beta_{l+j})_{C^{[d]}} = (\alpha_{l+i}, \alpha_{l+j})_C$ and $(\alpha_i, \alpha_j)_{C^{[d]}} = (\alpha_i, \alpha_j)_C$ for all i, j. In this case, we call $\beta_{l+1}, \cdots, \beta_{l+d}$ *added roots*; if β_i corresponds to $\alpha_i \in \dot{R}$ (resp. $\alpha_i \in \dot{\Delta}$), we call β_i an *added single root* (resp. *added single simple root*); if β_i corresponds to $\alpha_i \in 2\dot{R}$ (resp. $\alpha_i \in 2\dot{\Delta}$), we call β_i an *added double root* (resp. *added double simple root*).

Let C be a Cartan matrix, and let $(-, -)_C$ be the induced symmetric bilinear form of C. Then, given a root $\alpha \in \dot{R}$ we have $(\alpha, \alpha)_C = (\alpha_i, \alpha_i)_C$ for some simple root α_i. We call α a *short* (resp. *long*) *root* if $(\alpha, \alpha) = \min_i (\alpha_i, \alpha_i)$ (resp. $(\alpha, \alpha) = \max_i (\alpha_i, \alpha_i)$). Note that if C is symmetric, then all roots have the same square length, which we also call *long roots* by convention. If C is not symmetric, then every root is either short or long.

Accordingly, β_{l+i} is called an added single long (resp. short) root if β_{l+i} corresponds to a long (resp. short) root $\alpha_{l+i} \in \dot{R}$.

Definition 3.4. Let C be an indecomposable Cartan matrix of type X_l ($X = A, B, C, D, E, F, G$), and $\dot{R}$ be the root system of C. Let $C^{[d]}$ be a d-fold affinization matrix of C. Let b (resp. s) be the number of added single long (resp. short) roots, and let t be the number of added double roots. Then we call $C^{[d]}$ the d-fold affinization matrix of type $X_l^{(b,s,t)}$ ($b+s+t=d$).

Lemma 3.5. *Let C be an $l \times l$ Cartan matrix, and $A = (A_{ij})_{i,j\in I}$ be an $(l+d)$-matrix of rank l. Suppose that there is a subset $J \subset I$ such that $(A_{ij})_{i,j\in J} = C$. Then A is a d-fold affinization matrix $C^{[d]}$ if and only if for all $s \in I \setminus J$, $(A_{ij})_{i,j\in J\cup\{s\}}$ is a 1-fold affinization matrix $C^{[1]}$.*

Proof. The necessity is obvious. It remains to prove the sufficiency.

For all $s \in I \setminus J$, assume that the added root of the 1-fold affinization matrix $(A_{ij})_{i,j\in J\cup\{s\}}$ corresponds to $\alpha_s \in \dot{R} \cup 2\dot{R}$. Let $C^{[d]}$ be a d-fold affinization matrix of C, such that the added roots $\beta_{l+1}, \cdots, \beta_{l+d}$ correspond respectively to $\alpha_{l+1}, \cdots, \alpha_{l+d}$. Then for any $i \in J$ or $j \in J$ we have $A_{ij} = C^{[d]}_{ij}$, that is, after some suitable permutations of elements of the index set I, if the matrix A has a block form:

$$\begin{pmatrix} C & A_1 \\ A_2 & A_3 \end{pmatrix}$$

then the matrix $C^{[d]}$ has a corresponding block form:

$$\begin{pmatrix} C & A_1 \\ A_2 & A_3' \end{pmatrix}$$

Since both of them are of rank l, we have $A_3 = A_2C^{-1}A_1 = A_3'$. So $A = C^{[d]}$. □

Proposition 3.6. *Let C be a Cartan matrix. Then any d-fold affinization matrix $C^{[d]}$ of C is braid-equivalent to a standard d-fold affinization matrix of the same type.*

Proof. Let the root basis of C be $\dot{\Delta} = \{\alpha_1, \cdots, \alpha_l\}$, and the SIM-root basis of $C^{[d]}$ be $\Delta = \{\alpha_1, \cdots, \alpha_l, \beta_{l+1}, \cdots, \beta_{l+d}\}$, where $\beta_{l+1}, \cdots, \beta_{l+d}$ are added roots, corresponding respectively to $\alpha_{l+1}, \cdots, \alpha_{l+d} \in \dot{R} \cup 2\dot{R}$. Let $\alpha_{l+1} = s_{\alpha_{i_t}} \cdots s_{\alpha_{i_1}}(\alpha)$, where $\alpha \in \dot{\Delta} \cup 2\dot{\Delta}$, $\alpha_{i_j} \in \dot{\Delta}$ and $s_{\alpha_{i_j}}$ is a reflection for $j = 1, 2, \cdots, t$. Denote $\Delta' = \tau_{i_t,l+1}\Delta$, then $\Delta' = \{\alpha_1, \cdots, \alpha_l, s_{\alpha_{i_t}}(\beta_{l+1}), \beta_{l+2}, \cdots, \beta_{l+d}\}$ is an SIM-root basis braid-equivalent to Δ. Note that $s_{\alpha_{i_t}}(\beta_{l+1}) = \beta_{l+1} - \frac{2(\alpha_{i_t}, \alpha_{l+1})_C}{(\alpha_{i_t}, \alpha_{i_t})_C}\alpha_{i_t}$. It is easy to see that the structural matrix of Δ' is also a d-fold affinization and the added roots $s_{\alpha_{i_t}}(\beta_{l+1})$ correspond to $s_{\alpha_{i_t}}(\alpha_{l+1})$. By the induction on t we know that $C^{[d]}$ is braid-equivalent to a d-fold affinization matrix, and its SIM-root basis is $\{\alpha_1, \cdots, \alpha_l, \gamma_{l+1}, \beta_{l+2}, \cdots, \beta_{l+d}\}$, such that the added root γ_{l+1} corresponds to $\alpha \in \dot{\Delta} \cup 2\dot{\Delta}$. Applying the same argument to $\alpha_{l+2}, \cdots, \alpha_{l+d}$, we can show that $C^{[d]}$ is braid-equivalent to a d-fold affinization matrix, and its SIM-root basis is $\{\alpha_1, \cdots, \alpha_l, \gamma_{l+1}, \gamma_{l+2}, \cdots, \gamma_{l+d}\}$, such that the added roots $\gamma_{l+1}, \gamma_{l+2}, \cdots, \gamma_{l+d}$ correspond to the elements of $\dot{\Delta} \cup 2\dot{\Delta}$, that is, $C^{[d]}$ is braid-equivalent to a standard d-fold affinization matrix. □

Proposition 3.7. *Let $A = (A_{ij})_{i,j\in I}$ be an indecomposable generalized Cartan matrix of affine type. Then A is a 1-fold affinization matrix, and the type of A as a generalized Cartan matrix of affine type (see the diagrams in* [Kac]*) has the*

following relation with the type of A as a 1-fold affinization matrix:
(1) *A is of type $D_4^{(3)}$ if and only if A is of type $G_2^{(0,1,0)}$;*
(2) *A is of type $G_2^{(1)}$ if and only if A is of type $G_2^{(1,0,0)}$;*
(3) *A is of type $E_6^{(2)}$ if and only if A is of type $F_4^{(0,1,0)}$;*
(4) *A is of type $F_4^{(1)}$ if and only if A is of type $F_4^{(1,0,0)}$;*
(5) *A is of type $D_{l+1}^{(2)}$ if and only if A is of type $B_l^{(0,1,0)}$;*
(6) *A is of type $B_l^{(1)}$ if and only if A is of type $B_l^{(1,0,0)}$;*
(7) *A is of type $A_{2l}^{(2)}$ if and only if A is of type $B_l^{(0,0,1)}$;*
(8) *A is of type $A_{2l-1}^{(2)}$ if and only if A is of type $C_l^{(0,1,0)}$;*
(9) *A is of type $C_l^{(1)}$ if and only if A is of type $C_l^{(1,0,0)}$;*
(10) *A is of type $A_l^{(1)}$ if and only if A is of type $A_l^{(1,0,0)}$;*
(11) *A is of type $A_2^{(2)}$ if and only if A is of type $A_1^{(0,0,1)}$;*
(12) *A is of type $D_l^{(1)}$ if and only if A is of type $D_l^{(1,0,0)}$;*
(13) *A is of type $E_l^{(1)}$ if and only if A is of type $E_l^{(1,0,0)}$.*

Proof. Suppose A is of type $D_4^{(3)}$. Obviously A has an indecomposable Cartan submatrix $(A_{ij})_{i,j\in J}$ of type G_2, where J is a subset of I with $|I|-1$ elements. Denote $C=(A_{ij})_{i,j\in J}$ and let $C^{[1]}$ be a 1-fold affinization matrix of type $G_2^{(0,1,0)}$. By Theorem 2.10, $C^{[1]}$ is braid-equivalent to an indecomposable affine generalized Cartan matrix $A'=(A'_{ij})_{i,j\in I}$ such that the submatrix $(A'_{ij})_{i,j\in J}=C$. Note that for the 1-fold affinization matrix $C^{[1]}$, the added root corresponds to a short root of C. So the indecomposable affine generalized Cartan matrix A' has an indecomposable Cartan submatrix C of type G_2 and another simple root of A' (which is not a simple root of C) is a short root. Therefore the type of A' must be $D_4^{(3)}$, that is, $A'=A$. Hence A is a 1-fold affinization matrix and the statement (1) is true. Similarly for other cases. □

Definition 3.8. Let $A'=(A'_{ij})_{i,j\in I}$ be a positive semi-definite SIM, and $J\subsetneqq I$. We call $A'=(A'_{ij})_{i,j\in I}$ a *J-normal affinization matrix*, if $(A'_{ij})_{i,j\in J}$ is a Cartan matrix, and for any $s\in I\setminus J$, $(A'_{ij})_{i,j\in J\cup\{s\}}$ is a positive semi-definite generalized Cartan matrix of corank 1.

Let $A=(A_{ij})_{i,j\in I}$ be a positive semi-definite SIM. If A is braid-equivalent to a J-normal affinization matrix $A'=(A'_{ij})_{i,j\in I}$, then we say $A'=(A'_{ij})_{i,j\in I}$ is a *J-normal affinization matrix of A*.

Proposition 3.9. *Let $A=(A_{ij})_{i,j\in I}$ be a positive semi-definite symmetrizable intersection matrix with $\operatorname{rank}A=l<|I|$. For any subset $J\subset I$, if $|J|=l$ and $\operatorname{rank}(A_{ij})_{i,j\in J}=l$, then A is braid-equivalent to a J-normal affinization matrix under a sequence of APR-transformations at the vertices of J.*

Proof. Take $s_1\in I\setminus J$, then $(A_{ij})_{i,j\in J\cup\{s_1\}}$ is positive semi-definite of corank 1. By Theorem 2.10, under a sequence of APR-transformations at the vertices of J, A is braid-equivalent to a symmetrizable intersection matrix $A'=(A'_{ij})_{i,j\in I}$, such that $(A'_{ij})_{i,j\in J}$ is a Cartan matrix and $(A'_{ij})_{i,j\in J\cup\{s_1\}}$ is a positive semi-definite generalized Cartan matrix of corank 1. Take $s_2\in I\setminus(J\cup\{s_1\})$, then similarly under a sequence of APR-transformations at the vertices of J, A' is braid-equivalent to a symmetrizable intersection matrix $A''=(A''_{ij})_{i,j\in I}$, such that $(A''_{ij})_{i,j\in J}$ is a Cartan matrix and $(A''_{ij})_{i,j\in J\cup\{s_2\}}$ is a positive semi-definite generalized Cartan matrix of corank 1. Note that any symmetrizable generalized Cartan matrix is invariant under APR-transformations. So we have $(A''_{ij})_{i,j\in J\cup\{s_1\}} = (A'_{ij})_{i,j\in J\cup\{s_1\}}$. Now the induction on $|I\setminus J|$ implies the result. ☐

Lemma 3.10. *Let $A=(A_{ij})_{i,j\in I}$ be an indecomposable positive semi-definite symmetrizable intersection matrix of* $\operatorname{rank} A = l$. *Then there exists an indecomposable submatrix $(A_{ij})_{i,j\in J}$ such that $|J|=l$ and* $\operatorname{rank}(A_{ij})_{i,j\in J}=l$.

Proof. Since A is positive semi-definite and $\operatorname{rank} A = l$, there is a subset $K\subseteq I$ such that $|K|=l$ and$(A_{ij})_{i,j\in K}$ is a positive definite submatrix. Among all of these subsets K, we pick out a subset J, such that $(A_{ij})_{i,j\in J}$ has a minimal number of indecomposable components. We claim that $(A_{ij})_{i,j\in J}$ is indecomposable.

Otherwise, suppose $(A_{ij})_{i,j\in J}$ is decomposable. Let $(A_{ij})_{i,j\in J_1},\cdots,(A_{ij})_{i,j\in J_m}$ $(m>1)$ be all indecomposable components of $(A_{ij})_{i,j\in J}$. By Proposition 3.9, A is braid-equivalent to a J-normal affinization matrix $A'=(A'_{ij})_{i,j\in I}$. Obviously, $(A'_{ij})_{i,j\in J_1},\cdots,(A'_{ij})_{i,j\in J_m}$ are all indecomposable components of the Cartan submatrix $(A'_{ij})_{i,j\in J}$, and for all $s\in I\setminus J$, every indecomposable component of $(A'_{ij})_{i,j\in J\cup\{s\}}$ is of form $(A'_{ij})_{i,j\in J_{k_1}\cup\cdots\cup J_{k_n}\cup\{s\}}$, $(A'_{ij})_{i,j\in J_{k_{n+1}}},\cdots,(A'_{ij})_{i,j\in J_m}$, where $1\le n\le m$, and we also have $(A'_{ij})_{i,j\in J_{k_1}\cup\cdots\cup J_{k_n}\cup\{s\}}$ is an indecomposable affine generalized Cartan matrix.

If there is an $s\in I\setminus J$ such that the above $n>1$, that is, the number of all indecomposable components of $(A'_{ij})_{i,j\in J\cup\{s\}}$ is less that m, then we let $(A'_{ij})_{i,j\in J'}$ be the indecomposable Cartan submatrix of the indecomposable affine generalized Cartan matrix $(A'_{ij})_{i,j\in J_{k_1}\cup\cdots\cup J_{k_n}\cup\{s\}}$, such that $|J'|=|J_{k_1}\cup\cdots\cup J_{k_n}\cup\{s\}|-1$. Therefore $(A'_{ij})_{i,j\in J'\cup J_{k_{n+1}}\cup\cdots\cup J_{k_m}}$ is a Cartan matrix of order l, and the number of its indecomposable components is $1+(m-n)<m$, which contradicts to the choice of J. Therefore, for any $s\in I\setminus J$, if there is some $1\le x\le m$ such that $(A'_{ij})_{i,j\in J_{k_x}\cup\{s\}}$ is indecomposable, then $(A'_{ij})_{i,j\in J_{k_x}\cup\{s\}}$ must be an indecomposable affine generalized Cartan matrix.

Since A is indecomposable, A' is also indecomposable. Then there exist two submatrices among $(A'_{ij})_{i,j\in J_1},\cdots,(A'_{ij})_{i,j\in J_m}$, say $(A'_{ij})_{i,j\in J_1}$ and $(A'_{ij})_{i,j\in J_2}$, which are connected with each other by some vertices in $I\setminus J$, that is, there are $t_1,\cdots,t_k\in I$ $(k\ge 3)$ such that each of $A'_{t_1t_2},A'_{t_2t_3},\cdots,A'_{t_{k-1}t_k}$ is nonzero, where $t_1\in J_1$, $t_k\in J_2$, but $t_2,\cdots,t_{k-1}\in I\setminus J$. We can suppose that such k is minimal. Then for all $a\in J_1$, we have $A'_{at_3}=0$. By the result above we know that $(A'_{ij})_{i,j\in J_1\cup\{t_2\}}$ is an indecomposable affine generalized Cartan matrix. Then it is not difficult to show that $(A'_{ij})_{i,j\in J_1\cup\{t_2,t_3\}}$ is (or is braid-equivalent to) an indefinite generalized Cartan matrix, as contradicts to the positive semi-definiteness. ☐

In the following we come to one of our main results in this section, which shows that any indecomposable positive semi-definite symmetrizable intersection matrix is in fact braid-equivalent to a d-fold affinization matrix.

Theorem 3.11. *Let A be a positive semi-definite $(l+d)\times(l+d)$ symmetrizable intersection matrix with* $\operatorname{rank} A = l$. *Then A is braid-equivalent to a d-fold affinization matrix $C^{[d]}$.*

Proof. It is no harm to assume that A is indecomposable. We can choose some $J\subseteq I$ such that $|J|=l$ and $(A_{ij})_{i,j\in J}$ is an indecomposable symmetrizable intersection matrix with rank $(A_{ij})_{i,j\in J} = l$, according to Lemma 3.10. So by Proposition 3.9, A is braid-equivalent to a J-normal affine matrix $(A'_{ij})_{i,j\in I}$. Then the Cartan submatrix $(A'_{ij})_{i,j\in J}$ is also indecomposable , and for all $s \in I \setminus J$, $(A'_{ij})_{i,j\in J\cup\{s\}}$ is an indecomposable affine generalized Cartan submatrix. Now, if $(A'_{ij})_{i,j\in J}$ is of type X_l $(X = A, B, C, D, E, F, G)$, then we call J a *maximal subset* of A of type X_l. In the following we consider all possible cases and choose a suitable maximal subset J, and we prove the corresponding J-normal affinization matrix $(A'_{ij})_{i,j\in I}$ is a d-fold affinization matrix (for the concrete types of affine generalized Cartan matrices considered here, see the tables of the affine Dynkin diagrams in [Kac]).

(1) Suppose A has a maximal subset J of type G_2. Denote $(A'_{ij})_{i,j\in I}$ a J-normal affinization matrix of A. Then for all $s \in I\setminus J$, the type of the indecomposable affine generalized Cartan submatrix $(A'_{ij})_{i,j\in J\cup\{s\}}$ must be $G_2^{(1)}$ or $D_4^{(3)}$. We denote the number of $G_2^{(1)}$'s or $D_4^{(3)}$'s respectively by b or s. By Proposition 3.7 and Lemma 3.5, we know $(A'_{ij})_{i,j\in I}$ is a d-fold affinization matrix of type $G_2^{(b,s,0)}$.

(2) Suppose A has a maximal subset J of type F_4. Denote $(A'_{ij})_{i,j\in I}$ a J-normal affinization matrix of A. Then for any $s \in I \setminus J$, the type of the indecomposable affine generalized Cartan submatrix $(A'_{ij})_{i,j\in J\cup\{s\}}$ must be $F_4^{(1)}$ or $E_6^{(2)}$. We denote the number of $F_4^{(1)}$'s or $E_6^{(2)}$'s respectively by b or s. By Proposition 3.7 and Lemma 3.5, we know $(A'_{ij})_{i,j\in I}$ is a d-fold affinization matrix of type $F_4^{(b,s,0)}$.

(3) Suppose A has a maximal subset J of type B_l $(l \geq 3)$ and has no maximal subset of type F_4. Denote $(A'_{ij})_{i,j\in I}$ a J-normal affinization matrix of A. Then for all $s \in I \setminus J$, the type of the indecomposable affine generalized Cartan submatrix $(A'_{ij})_{i,j\in J\cup\{s\}}$ must be $B_l^{(1)}$, $D_{l+1}^{(2)}$ or $A_{2l}^{(2)}$. Denote the number of them by b, s or t respectively. By Proposition 3.7 and Lemma 3.5, we know that $(A'_{ij})_{i,j\in I}$ is a d-fold affinization matrix of type $B_l^{(b,s,t)}$.

(4) Suppose A has a maximal subset J of type B_2. Let $(A'_{ij})_{i,j\in I}$ be a J-normal affinization matrix of A. Then for all $s \in I\setminus J$, the type of the indecomposable affine generalized Cartan submatrix $(A'_{ij})_{i,j\in J\cup\{s\}}$ must be $C_2^{(1)}$, $D_3^{(2)}$ or $A_4^{(2)}$. Denote the number of them by b, s or t respectively. By Proposition 3.7 and Lemma 3.5, we know that $(A'_{ij})_{i,j\in I}$ is a d-fold affinization matrix of type $B_2^{(b,s,t)}$.

(5) Suppose A has a maximal subset J of type C_l $(l \geq 3)$ and has no maximal subset of type F_4 or B_l. Let $(A'_{ij})_{i,j\in I}$ be a J-normal affinization matrix of A. Then for any $s \in I \setminus J$, the type of the indecomposable affine generalized Cartan submatrix $(A'_{ij})_{i,j\in J\cup\{s\}}$ must be $C_l^{(1)}$ or $A_{2l-1}^{(2)}$. Denote the number of them by b or s respectively. By Proposition 3.7 and Lemma 3.5, we know that $(A'_{ij})_{i,j\in I}$ is a d-fold affinization matrix of type $C_l^{(b,s,0)}$.

(6) Suppose A is not symmetric and has no maximal subset of type G_2, F_4, B_l or C_l. It is easy to see rank $A = 1$. Take any $j \in I$ such that the length of the root α_j in the corresponding SIM-root basis is minimal. Denote $J = \{j\}$ and let $(A'_{ij})_{i,j\in I}$ be a J-normal affinization matrix of A. Then for any $s \in I \setminus J$, the type of the indecomposable affine generalized Cartan submatrix $(A'_{ij})_{i,j\in J\cup\{s\}}$ must be $A_1^{(1)}$ or $A_2^{(2)}$. Denote the number of them by b or t respectively. By Proposition 3.7 and Lemma 3.5, we know that $(A'_{ij})_{i,j\in I}$ is a d-fold affinization matrix of type $A_1^{(b,0,t)}$.

We have discussed the case that A is not symmetric as above. In the following, we consider the case that A is symmetric.

(7) Suppose A has a maximal subset J of type E_l $(l = 6, 7, 8)$. Denote $(A'_{ij})_{i,j\in I}$ a J-normal affinization matrix of A. Then for any $s \in I \setminus J$, the type of the indecomposable affine generalized Cartan submatrix $(A'_{ij})_{i,j\in J\cup\{s\}}$ must be $E_l^{(1)}$. By Proposition 3.7 and Lemma 3.5, we know $(A'_{ij})_{i,j\in I}$ is a d-fold affinization matrix of type $E_l^{(d,0,0)}$.

(8) Suppose A has a maximal subset J of type D_l and has no maximal subset of type E_l. Denote $(A'_{ij})_{i,j\in I}$ a J-normal affinization matrix of A. Then for any $s \in I \setminus J$, the type of the indecomposable affine generalized Cartan submatrix $(A'_{ij})_{i,j\in J\cup\{s\}}$ must be $D_l^{(1)}$. By Proposition 3.7 and Lemma 3.5, we know $(A'_{ij})_{i,j\in I}$ is a d-fold affinization matrix of type $D_l^{(d,0,0)}$.

(9) Suppose A has a maximal subset J of type A_l and has no maximal subset of other types. Denote $(A'_{ij})_{i,j\in I}$ a J-normal affinization matrix of A. Then for any $s \in I \setminus J$, the type of the indecomposable affine generalized Cartan submatrix $(A'_{ij})_{i,j\in J\cup\{s\}}$ must be $A_l^{(1)}$. By Proposition 3.7 and Lemma 3.5, we know $(A'_{ij})_{i,j\in I}$ is a d-fold affinization matrix of type $A_l^{(d,0,0)}$.

We have discussed all possible cases, and in each case, A is braid-equivalent to a d-fold affinization matrix which is at the same time a normal affinization matrix. ☐

3.2. **Classification of Positive Semi-definite Symmetrizable Intersection Matrices.**

As another main result of this section, the following theorem produces a classification of all indecomposable positive semi-definite symmetrizable intersection matrices by standard d-fold affinization matrices, up to braid-equivalence.

Theorem 3.12. *Let A be an indecomposable positive semi-definite symmetrizable intersection matrix. Then A is braid-equivalent to a standard d-fold affinization matrix of type $X_l^{(b,s,t)}$ $(b+s+t = d)$. Moreover, any two standard d-fold affinization matrices with the same type are braid-equivalent. Here the type $X_l^{(b,s,t)}$ must be one of the following forms:*
(1) $X_l^{(d,0,0)}$ *for* $X_l = A_l (l \geq 2)$, D_l, E_6, E_7, E_8;
(2) $X_l^{(b,s,0)}$ *for* $X_l = C_l (l \geq 3)$, F_4, G_2;
(3) $A_1^{(b,0,t)}$, $B_l^{(b,s,t)}$, *where for* $B_l^{(b,s,t)}$, *any added double root must correspond to the double of a short root.*

Proof. By Theorem 3.11 and Proposition 3.6, A is braid-equivalent to a standard d-fold affinization matrix.

If $A' = (A'_{ij})_{i,j\in I}$ and $A'' = (A''_{ij})_{i,j\in I}$ are two standard d-fold affinization matrices of the same type $X_l^{(b,s,t)}$, then we can assume that there is a subset $J \subseteq I$ such that $(A'_{ij})_{i,j\in J} = (A''_{ij})_{i,j\in J}$ is an indecomposable Cartan matrix of type X_l and for any $a \in J \setminus I$, $(A'_{ij})_{i,j\in J\cup\{a\}}$ and $(A''_{ij})_{i,j\in J\cup\{a\}}$ are two standard 1-fold affinization matrices of the same type. So for all $a \in J \setminus I$, by Proposition 3.7 we see both $(A'_{ij})_{i,j\in J\cup\{a\}}$ and $(A''_{ij})_{i,j\in J\cup\{a\}}$ are braid-equivalent to the same one indecomposable affine generalized Cartan matrix. Therefore both A' and A'' are braid-equivalent to the same one J-normal affinization matrix. Hence they are braid-equivalent to each other.

It is easy to see that there exist standard d-fold affinization matrices of type appearing in (1), (2) and (3). Next we show that the types appearing in (1), (2) and (3) are the only possibilities. For (1) and (2), we only need to prove the added roots in a standard d-fold affinization matrix $C^{[d]}$ are all single roots, that is, the corresponding roots $\alpha_{l+1}, \cdots, \alpha_{l+d}$ lie in $\dot{\Delta}$. Otherwise, suppose some $\alpha_{l+j} = 2\alpha_k \in 2\dot{\Delta}$, $1 \leq k \leq l$. Then for all $i = 1, \cdots, l$,

$$C^{[d]}_{l+j,\ i} = \frac{2(\alpha_{l+j},\ \alpha_i)_C}{(\alpha_{l+j},\ \alpha_{l+j})_C} = \frac{(\alpha_k,\ \alpha_i)_C}{(\alpha_k,\ \alpha_k)_C} = \frac{1}{2} \cdot C_{ki}.$$

Obviously, for any Cartan matrix C of type $A_l (l \geq 2)$, $C_l (l \geq 3)$, D_l, E_6, E_7, E_8, F_4 or G_2, there exists an i with $1 \leq i \leq l$ such that $C_{ki} = -1$ or -3. Hence $C^{[d]}_{l+1,\ i} = \frac{1}{2} \cdot C_{ki} \notin \mathbf{Z}$, as contradicts to the definition of d-fold affinizations.

For (3), we only need to prove that for the type $B_l^{(b,s,t)}$, any added double root must correspond to the double of a short root. Let $\{\alpha_1, \cdots, \alpha_{l-1}, \alpha_l\}$ be a root basis of B_l, where $\alpha_1, \cdots, \alpha_{l-1}$ are the long simple roots and α_l is the short simple root. Let $C^{[d]}$ be a standard d-affinization matrix of type $B_l^{(b,s,t)}$. Suppose β_{l+j} is an added double root corresponding to $\alpha_{l+j} = 2\alpha_k$ with $1 \leq k \leq l-1$, then

$$C^{[d]}_{l+j,\ k+1} = \frac{2(\alpha_{l+j},\ \alpha_{k+1})_C}{(\alpha_{l+j},\ \alpha_{l+j})_C} = \frac{(\alpha_k,\ \alpha_{k+1})_C}{(\alpha_k,\ \alpha_k)_C} = \frac{1}{2} \cdot C_{k,\ k+1} = -\frac{1}{2}.$$

This is impossible and so any added double root must correspond to the double of a short root. $\square$

4. Weyl Root Systems of Positive Semi-definite SIM's

In this section, we give an explicit structure of the Weyl root system for each d-fold affinization matrices by using the root system of the Cartan matrix and some special null roots.

Let A be an SIM, and $(-,-)$ be the corresponding symmetric bilinear form. Set

$$\mathrm{rad}(-,-) := \{\alpha \in H \mid (\alpha,\ \beta) = 0 \quad \text{for all} \quad \beta \in H\}$$

and

$$\mathrm{rad}(\Gamma) := \mathrm{rad}(-,-) \cap \Gamma,$$

where Γ is the root lattice. We call each element in $\mathrm{rad}(\Gamma)$ a *null root*.

For the sake of convenience, we relabel the simple roots in Dynkin diagrams as follows:

A_l:



B_l $(l \geq 3)$:



C_l:



D_l:



E_6:



E_7:



E_8:



F_4:



G_2:



In the above labeling , α_1 is always a long root and α_l(where l is the maximal number in the labeling) is always a short root.

Let A be an SIM which is positive semi-definite. To investigate its Weyl root system R^W, we can assume that A is a standard d-fold affinization matrix of type $X_l^{(b,s,t)}$. We can further assume that A is so 'standard' that each added long root corresponds to α_1, each added short root corresponds to α_l, and each added double short root corresponds to $2\alpha_l$.

Therefore we can assume that all added long roots corresponding to α_1 are $\alpha_{1j}, j = 1, 2, \cdots, b$; all added short roots corresponding to α_l are $\alpha_{lj}, j = 1, 2, \cdots, s$; and all added double short roots corresponding to $2\alpha_l$ are $\beta_{lj}, j = 1, 2, \cdots, t$. Denote $\delta_{1j} := \alpha_{1j} - \alpha_1$, $j = 1, \cdots, b$, $\delta_{lj} := \alpha_{lj} - \alpha_l$, $j = 1, \cdots, s$, and $\delta'_{lj} := \beta_{lj} - 2\alpha_l$, $j = 1, \cdots, t$. It is easy to see that $\{\delta_{11}, \cdots, \delta_{1b}, \delta_{l1}, \cdots, \delta_{ls}, \delta'_{l1}, \cdots, \delta'_{lt}\}$ is a basis of rad$(-,-)$.

By the classification theorem (Theorem 3.12) we know that the form of the type $X_l^{(b,s,t)}$ is: (1) $X_l^{(d,0,0)}$, where $X_l = A_l (l \geq 2)$, D_l, E_6, E_7, E_8; (2) $X_l^{(b,s,0)}$, where $X_l = C_l (l \geq 3)$, F_4, G_2; or (3) $A_1^{(b,0,t)}$, $B_l^{(b,s,t)}$. Thus the following theorem gives a complete description of the Weyl root system R^W of A in terms of the root system $\dot{R}$ of the Cartan matrix of type X_l and the null roots $\delta_{11}, \cdots, \delta_{1b}, \delta_{l1}, \cdots, \delta_{ls}, \delta'_{l1}, \cdots, \delta'_{lt}$. Note that these Weyl roots have at most three lengths. So we denote all short roots, all middle roots and all long roots respectively by R_s^W, R_m^W and R_l^W.

Theorem 4.1. *The assumptions are as above. Then*

(1) *For the type $X_l^{(d,0,0)}$, where $X_l = A_l (l \geq 2)$, D_l, E_6, E_7, E_8, we have*

$$R^W = \dot{R} + rad(\Gamma).$$

(2) *For the type $X_l^{(b,s,0)}$, where $X_l = C_l (l \geq 3)$, F_4, G_2, we have*

$$R^W = R_s^W \bigcup R_l^W,$$

where

$$R_s^W = \dot{R}_s + rad(\Gamma)$$

and R_l^W is the following set:

(2.1) *for $X_l = C_l (l \geq 3)$,*

$$\begin{aligned} R_l^W \quad = \quad & \{\alpha + \sum_{j=1}^{b} a_j \delta_{1j} + 2 \sum_{j=1}^{s} b_j \delta_{lj} \mid \alpha \in \dot{R}_l, a_j, b_j \in \mathbf{Z}, \\ & \text{and there is at most one odd } a_j, j = 1, \cdots, b\}; \end{aligned}$$

(2.2) *for $X_l = F_4$,*

$$R_l^W = \{\alpha + \sum_{j=1}^{b} a_j \delta_{1j} + 2 \sum_{j=1}^{s} b_j \delta_{lj} \mid \alpha \in \dot{R}_l, a_j, b_j \in \mathbf{Z}\};$$

(2.3) *for $X_l = G_2$,*

$$R_l^W = \{\alpha + \sum_{j=1}^{b} a_j \delta_{1j} + 3 \sum_{j=1}^{s} b_j \delta_{lj} \mid \alpha \in \dot{R}_l, a_j, b_j \in \mathbf{Z}\}.$$

(3) *For the type $A_1^{(b,0,t)}$, we have $R^W = R_s^W \bigcup R_l^W$, where*

$$\begin{aligned} R_s^W \quad = \quad & \{\pm\alpha_1 + \sum_{j=1}^{b} a_j \delta_{1j} + \sum_{j=1}^{t} b_j \delta'_{1j} \mid a_j, b_j \in \mathbf{Z}, \\ & \text{and there is at most one odd } a_j, j = 1 \cdots, b\}. \end{aligned}$$

and

$$\begin{aligned} R_l^W \quad = \quad & \{\pm 2\alpha_1 + 4 \sum_{j=1}^{b} a_j \delta_{1j} + \sum_{j=1}^{t} b_j \delta'_{1j} \mid a_j, b_j \in \mathbf{Z}, \\ & \text{and there is exactly one odd } b_j, j = 1, \cdots, t\}. \end{aligned}$$

(4) *For the type $B_l^{(b,s,t)}$, we have $R^W = R_s^W \bigcup R_m^W \bigcup R_l^W$, where*

$$\begin{aligned} R_s^W \quad = \quad & \{\alpha + \sum_{j=1}^{b} a_j \delta_{1j} + \sum_{j=1}^{s} b_j \delta_{lj} + \sum_{j=1}^{t} c_j \delta'_{lj} \mid \alpha \in \dot{R}_s, a_j, b_j, c_j \in \mathbf{Z}, \\ & \text{and there is at most one odd } b_j, j = 1, \cdots, s\}; \end{aligned}$$

$$\begin{aligned} R_l^W &= \{2\alpha + 2\sum_{j=1}^{b} a_j\delta_{1j} + 4\sum_{j=1}^{s} b_j\delta_{lj} + \sum_{j=1}^{t} c_j\delta'_{lj} \mid \alpha \in \dot{R}_s, a_j, b_j, c_j \in \mathbf{Z}, \\ &\qquad \textit{and there is exactly one odd } c_j, j = 1, \cdots, t\}. \end{aligned}$$

and R_m^W *is the following set:*

(4.1) *for* $l = 2$,

$$\begin{aligned} R_m^W &= \{\alpha + \sum_{j=1}^{b} a_j\delta_{1j} + 2\sum_{j=1}^{s} b_j\delta_{lj} + \sum_{j=1}^{t} c_j\delta'_{lj} \mid \alpha \in \dot{R}_l, a_j, b_j, c_j \in \mathbf{Z}, \\ &\qquad \textit{and there is at most one odd } a_j, j = 1 \cdots, b\}; \end{aligned}$$

(4.2) *for* $l > 2$,

$$R_m^W = \{\alpha + \sum_{j=1}^{b} a_j\delta_{1j} + 2\sum_{j=1}^{s} b_j\delta_{lj} + \sum_{j=1}^{t} c_j\delta'_{lj} \mid \alpha \in \dot{R}_l, a_j, b_j, c_j \in \mathbf{Z}\}.$$

Before proving this theorem, we first show the following lemmas.

Lemma 4.2. *If* $\Delta \subseteq \dot{R} + rad(\Gamma)$, *then* $R_s^W \subseteq \dot{R}_s + rad(\Gamma)$ *and* $R_l^W \subseteq \dot{R}_l + rad(\Gamma)$.

Proof. For $\alpha, \beta \in \dot{R} + rad(\Gamma)$, let $\alpha = \dot{\alpha} + \delta$, $\beta = \dot{\beta} + \delta'$, where $\dot{\alpha}, \dot{\beta} \in \dot{R}$ and $\delta, \delta' \in rad(\Gamma)$. Then $s_\alpha(\beta) = \beta - (\alpha^\vee, \beta)\alpha = \dot{\beta} + \delta' - (\alpha^\vee, \beta)(\dot{\alpha} + \delta) = \dot{\beta} + \delta' - (\dot{\alpha}^\vee, \dot{\beta})(\dot{\alpha} + \delta) = s_{\dot{\alpha}}(\dot{\beta}) + (\delta' - (\dot{\alpha}^\vee, \dot{\beta})\delta) \in \dot{R} + rad(\Gamma)$. Therefore if $\beta \in \dot{R}_s + rad(\Gamma)$, then $s_\alpha(\beta) \in \dot{R}_s + rad(\Gamma)$; if $\beta \in \dot{R}_l + rad(\Gamma)$, then $s_\alpha(\beta) \in \dot{R}_l + rad(\Gamma)$. So by induction on the number of reflections, we have $R_s^W \subseteq \dot{R}_s + rad(\Gamma)$ and $R_l^W \subseteq \dot{R}_l + rad(\Gamma)$. ☐

Lemma 4.3. *If* $\alpha \in R^W$ *and* $\delta \in rad(\Gamma)$, *then* $\alpha + \delta \in R^W$ *if and only if* $\alpha - \delta \in R^W$.

Proof. The result follows since $\alpha - \delta = -s_\alpha(\alpha + \delta)$. ☐

Lemma 4.4. *For* $\alpha, \beta \in R^W$ *with* $(\alpha^\vee, \beta) = -1$ *and* $\delta \in rad(\Gamma)$, *if* $\alpha + \delta \in R^W$, *then* $\beta + \delta \in R^W$. *As a consequence, for any* $\alpha \in \dot{R}$, *we have* $\alpha + \delta_{1j} \in R^W$ *for* $j = 1, \cdots, b$.

Proof. Clearly $\beta + \delta = s_\alpha s_{\alpha+\delta}(\beta) \in R^W$. Furthermore, note that $\alpha_1 + \delta_{1j} = \alpha_{1j} \in R^W$ for $j = 1, \cdots, b$. For any $\alpha_i, i = 2, \cdots, l$, there is an $\alpha_k, k < i$, such that $(\alpha_k^\vee, \alpha_i) = -1$. By induction, we have $\alpha_i + \delta_{1j} \in R^W$ for $i = 1, \cdots, l$ and $j = 1, \cdots, b$. Therefore, for any $\alpha \in \dot{R}$, we have $\alpha + \delta_{1j} \in R^W$ for $j = 1, \cdots, b$. ☐

Lemma 4.5. (1) *For* $C_l (l \geq 2)$, *if* $\alpha \in \dot{R}$ *and* $\beta \in \dot{R}_l$, *then* $(\alpha^\vee, \beta)$ *is even;*
(2) *for* $B_l (l \geq 3)$, *if* $\alpha \in \dot{R}_s$ *and* $\beta \in \dot{R}$, *then* $(\alpha^\vee, \beta)$ *is even;*
(3) *for* F_4, *if* $\alpha \in \dot{R}_s$ *and* $\beta \in \dot{R}_l$, *then* $(\alpha^\vee, \beta)$ *is even;*
(4) *for* G_2, *if* $\alpha \in \dot{R}_s$ *and* $\beta \in \dot{R}_l$, *then* $(\alpha^\vee, \beta)$ *is divided by 3.*

Proof. In all cases, when α, β are simple roots, the result is true. Note that

$$\begin{aligned} ((s_{\gamma_1}(\alpha))^\vee, s_{\gamma_2}(\beta)) &= (\alpha^\vee, \beta) - (\alpha^\vee, \gamma_1)(\gamma_1^\vee, \beta) - (\alpha^\vee, \gamma_2)(\gamma_2^\vee, \beta) \\ &\quad + (\alpha^\vee, \gamma_1)(\gamma_1^\vee, \gamma_2)(\gamma_2^\vee, \beta). \end{aligned}$$

So by induction on the number of reflections, we obtain the result. ☐

Proof of theorem 4.1. (1) Consider the type $X_l^{(d,0,0)}$, where $X_l = A_l (l \geq 2)$, D_l, E_6, E_7, E_8. Since $\alpha_{1j} = \alpha_1 + \delta_{1j}$ for $j = 1, \cdots, s$, we have $\alpha \in \dot{R} + rad(\Gamma)$ for any $\alpha \in \Delta$. By Lemma 4.2, we obtain $R^W \subseteq \dot{R} + rad(\Gamma)$.

On the other hand, it suffices to show that $\{\alpha_i + \sum_{j=1}^{d} a_j \delta_{1j} \mid i = 1, 2, \cdots, l,\ a_j \in \mathbf{Z}\} \subseteq R^W$. For any $\alpha_i, i = 2, \cdots, l$, there is an $\alpha_k, k < i$, such that $(\alpha_k^\vee, \alpha_i) = -1$. So by Lemma 4.4, we only need to show that $S_1 := \{\alpha_1 + \sum_{j=1}^{d} a_j \delta_{1j} \mid a_j \in \mathbf{Z}\} \subseteq R^W$. Given any $\gamma \in S_1$, we have $s_{\alpha_2} s_{\alpha_2 + \delta_{1j}}(\gamma) = \gamma + \delta_{1j}$. Therefore if $\gamma \in R^W$, we have $\gamma \pm \delta_{1j} \in R^W$ for $j = 1, \cdots, d$. By induction on the number $n = \sum_{j=1}^{d} |a_j|$, we obtain that $S_1 \subseteq R^W$.

(2) Consider the type $X_l^{(b,s,0)}$, where $X_l = C_l (l \geq 3), F_4, G_2$. Since $\alpha_{lj} = \alpha_l + \delta_{lj}$ for $j = 1, \cdots, s$, we have $\Delta_s \subseteq \dot{R}_s + rad(\Gamma)$. By Lemma 4.2, we obtain $R_s^W \subseteq \dot{R}_s + rad(\Gamma)$.

On the other hand, by Lemma 4.4, we only need to prove that $S_2 := \{\alpha_l + \sum_{j=1}^{b} a_j \delta_{1j} + \sum_{j=1}^{s} b_j \delta_{lj} \mid a_j, b_j \in \mathbf{Z}\} \subseteq R_s^W$. By Lemma 4.4, we have $\alpha_i + \delta_{1j} \in R^W$ for $i = 1, \cdots, l$, and $j = 1, \cdots, b$. For $C_l (l \geq 3)$ and F_4, as $\alpha_l + \delta_{lj} = \alpha_{lj} \in R^W$ for $j = 1, \cdots, s$, by Lemma 4.4, we have $\alpha_{l-1} + \delta_{lj} \in R^W$ for $j = 1, \cdots, s$; for G_2, as $\alpha_2 + \delta_{2j} = \alpha_{2j} \in R^W$ for $j = 1, \cdots, s$, we have $\alpha_1 + 3\delta_{2j} = s_{\alpha_2} s_{\alpha_2 + \delta_{2j}}(\alpha_1) \in R^W$ for $j = 1, \cdots, s$. Given any $\gamma \in S_2$, we have $s_{\alpha_{l-1}} s_{\alpha_{l-1} + \delta_{1j}}(\gamma) = \gamma + \delta_{1j}$. Furthermore, for $C_l (l \geq 3)$ and F_4, we have $s_{\alpha_{l-1}} s_{\alpha_{l-1} + \delta_{lj}}(\gamma) = \gamma + \delta_{lj}$; for G_2 we have $s_{\alpha_1} s_{\alpha_1 + 3\delta_{2j}} s_{\alpha_2} s_{\alpha_2 + \delta_{2j}}(\gamma) = \gamma + \delta_{2j}$. So, if $\gamma \in R^W$, we have $\gamma \pm \delta_{1j}, \gamma \pm \delta_{2k} \in R^W$ for all $j = 1, \cdots, b$, and $k = 1, \cdots, s$. By induction on the number $n = \sum_{j=1}^{b} |a_j| + \sum_{j=1}^{s} |b_j|$, we obtain $S_2 \subseteq R_s^W$.

(2.1) $X_l = C_l (l \geq 3)$. Let S_3 be the set on the right-hand side. Since $\alpha_{1j} = \alpha_1 + \delta_{1j}$ for $j = 1, \cdots, b$, we have $\Delta_l \subseteq S_3$. Moreover, for any $\beta = \alpha + \sum_{j=1}^{b} a_j \delta_{1j} + 2 \sum_{j=1}^{s} b_j \delta_{lj} \in S_3$ with $\alpha \in \dot{R}_l$, we have

$$s_{\alpha_{1k}}(\beta) = s_{\alpha_{1k}}(\alpha) + \sum_{j=1}^{b} a_j \delta_{1j} + 2 \sum_{j=1}^{s} b_j \delta_{lj} = s_{\alpha_1}(\alpha) - (\alpha_1^\vee, \alpha)\delta_{1k} + \sum_{j=1}^{b} a_j \delta_{1j} + 2 \sum_{j=1}^{s} b_j \delta_{lj};$$

$$s_{\alpha_{lk}}(\beta) = s_{\alpha_{lk}}(\alpha) + \sum_{j=1}^{b} a_j \delta_{1j} + 2 \sum_{j=1}^{s} b_j \delta_{lj} = s_{\alpha_l}(\alpha) - (\alpha_l^\vee, \alpha)\delta_{lk} + \sum_{j=1}^{b} a_j \delta_{1j} + 2 \sum_{j=1}^{s} b_j \delta_{lj}.$$

By Lemma 4.5, both $(\alpha_1^\vee, \alpha)$ and $(\alpha_l^\vee, \alpha)$ are even. Then we have

$$s_{\alpha_{11}}(\beta), \cdots, s_{\alpha_{1b}}(\beta), s_{\alpha_{l1}}(\beta), \cdots, s_{\alpha_{ls}}(\beta) \in S_3,$$

and $s_{\alpha_i}(\beta) = s_{\alpha_i}(\alpha) + \sum_{j=1}^{b} a_j \delta_{1j} + 2 \sum_{j=1}^{s} b_j \delta_{lj} \in S_3$ for $i = 1, 2, \cdots, l$. So by induction on the number of reflections, we obtain $R_l^W \subseteq S_3$.

Next we prove $S_3 \subseteq R_l^W$. It suffices to show that $S_3' \subseteq R_l^W$, where $S_3' := \{\alpha_1 +$

$\sum_{j=1}^{b} a_j\delta_{1j} + 2\sum_{j=1}^{s} b_j\delta_{lj} \mid a_j, b_j \in \mathbf{Z}$, and there is at most one odd $a_j, j = 1, \cdots, b\}$. By Lemma 4.4, we have $\alpha_i + \delta_{1j} \in R^W$ for $i = 1, \cdots, l$ and $j = 1, \cdots, b$. And since $\alpha_l + \delta_{lj} = \alpha_{lj} \in R^W$ for $j = 1, \cdots, s$, by Lemma 4.4, we have $\alpha_i + \delta_{lj} \in R^W$ for $i = l-1, \cdots, 2$ and $j = 1, \cdots, s$. For any $\gamma \in S_3'$, we have $s_{\alpha_2}s_{\alpha_2+\delta_{1j}}(\gamma) = \gamma + 2\delta_{1j}$ and $s_{\alpha_2}s_{\alpha_2+\delta_{lj}}(\gamma) = \gamma + 2\delta_{lj}$. Therefore, if $\gamma \in R^W$, we have $\gamma \pm 2\delta_{1j}, \gamma \pm 2\delta_{lk} \in R^W$ for all $j = 1, \cdots, b$ and $k = 1, \cdots, s$. By induction on the number $n = \sum_{j=1}^{b} |a_j| + 2\sum_{j=1}^{s} |b_j|$, we obtain $S_3' \subseteq R_l^W$.

(2.2) $X_l = F_4$. Let S_4 be the set on the right-hand side. Since $\alpha_{1j} = \alpha_1 + \delta_{1j}$ for $j = 1, \cdots, b$, we have $\Delta_l \subseteq S_4$. Moreover, for any $\beta = \alpha + \sum_{j=1}^{b} a_j\delta_{1j} + 2\sum_{j=1}^{s} b_j\delta_{4j} \in S_4$ with $\alpha \in \dot{R}_l$, we have

$$s_{\alpha_{1k}}(\beta) = s_{\alpha_{1k}}(\alpha) + \sum_{j=1}^{b} a_j\delta_{1j} + 2\sum_{j=1}^{s} b_j\delta_{4j} = s_{\alpha_1}(\alpha) - (\alpha_1^\vee, \alpha)\delta_{1k} + \sum_{j=1}^{b} a_j\delta_{1j} + 2\sum_{j=1}^{s} b_j\delta_{4j};$$

$$s_{\alpha_{4k}}(\beta) = s_{\alpha_{4k}}(\alpha) + \sum_{j=1}^{b} a_j\delta_{1j} + 2\sum_{j=1}^{s} b_j\delta_{4j} = s_{\alpha_4}(\alpha) - (\alpha_4^\vee, \alpha)\delta_{4k} + \sum_{j=1}^{b} a_j\delta_{1j} + 2\sum_{j=1}^{s} b_j\delta_{4j}.$$

By Lemma 4.5, $(\alpha_4^\vee, \alpha)$ are even. Then we have

$$s_{\alpha_{11}}(\beta), \cdots, s_{\alpha_{1b}}(\beta), s_{\alpha_{41}}(\beta), \cdots, s_{\alpha_{4s}}(\beta) \in S_4,$$

and $s_{\alpha_i}(\beta) = s_{\alpha_i}(\alpha) + \sum_{j=1}^{b} a_j\delta_{1j} + 2\sum_{j=1}^{s} b_j\delta_{4j} \in S_4$ for $i = 1, 2, 3, 4$. So by induction on the number of reflections, we obtain $R_l^W \subseteq S_4$.

It remains to prove $S_4 \subseteq R_l^W$. By Lemma 4.4, it suffices to show $S_4' := \{\alpha_2 + \sum_{j=1}^{b} a_j\delta_{1j} + 2\sum_{j=1}^{s} b_j\delta_{4j} \mid a_j, b_j \in \mathbf{Z}\} \subseteq R_l^W$. Since $\alpha_4 + \delta_{4j} = \alpha_{4j} \in R^W$ for $j = 1, \cdots, s$, we have $\alpha_3 + \delta_{4j} \in R^W$ by Lemma 4.4. For any $\gamma \in S_4'$, we have $s_{\alpha_1}s_{\alpha_{1j}}(\gamma) = \gamma + \delta_{1j}$ and $s_{\alpha_3}s_{\alpha_3+\delta_{4j}}(\gamma) = \gamma + 2\delta_{4j}$. Thus, if $\gamma \in R^W$, we have $\gamma \pm \delta_{1j}, \gamma \pm 2\delta_{4k} \in R^W$ for all $j = 1, \cdots, b$ and $k = 1, \cdots, s$. By induction on the number $n = \sum_{j=1}^{b} |a_j| + 2\sum_{j=1}^{s} |b_j|$, we have $S_4' \subseteq R_l^W$.

(2.3) $X_l = G_2$. Let S_5 be the set on the right-hand side. Since $\alpha_{1j} = \alpha_1 + \delta_{1j}$ for $j = 1, \cdots, b$, we obtain $\Delta_l \subseteq S_5$. And for any $\beta = \alpha + \sum_{j=1}^{b} a_j\delta_{1j} + 3\sum_{j=1}^{s} b_j\delta_{2j} \in S_5$, where $\alpha \in \dot{R}_l$, we have

$$s_{\alpha_{1k}}(\beta) = s_{\alpha_{1k}}(\alpha) + \sum_{j=1}^{b} a_j\delta_{1j} + 3\sum_{j=1}^{s} b_j\delta_{2j} = s_{\alpha_1}(\alpha) - (\alpha_1^\vee, \alpha)\delta_{1k} + \sum_{j=1}^{b} a_j\delta_{1j} + 3\sum_{j=1}^{s} b_j\delta_{2j};$$

$$s_{\alpha_{2k}}(\beta) = s_{\alpha_{2k}}(\alpha) + \sum_{j=1}^{b} a_j\delta_{1j} + 3\sum_{j=1}^{s} b_j\delta_{2j} = s_{\alpha_2}(\alpha) - (\alpha_2^\vee, \alpha)\delta_{2k} + \sum_{j=1}^{b} a_j\delta_{1j} + 3\sum_{j=1}^{s} b_j\delta_{2j}.$$

By Lemma 4.5, $(\alpha_2^\vee, \alpha)$ is divided by 3. Then we obtain

$$s_{\alpha_{11}}(\beta), \cdots, s_{\alpha_{1b}}(\beta), s_{\alpha_{21}}(\beta), \cdots, s_{\alpha_{2s}}(\beta) \in S_5,$$

and $s_{\alpha_i}(\beta) = s_{\alpha_i}(\alpha) + \sum\limits_{j=1}^{b} a_j\delta_{1j} + 3\sum\limits_{j=1}^{s} b_j\delta_{2j} \in S_5$ for $i = 1, 2$. So by induction on the number of reflections, we get $R_l^W \subseteq S_5$.

Next we prove $S_5 \subseteq R_l^W$. We only need to show that $S_5' \subseteq R_l^W$, where $S_5' := \{\alpha_1 + \sum\limits_{j=1}^{b} a_j\delta_{1j} + 3\sum\limits_{j=1}^{s} b_j\delta_{2j} \mid a_j, b_j \in \mathbf{Z}\}$. By Lemma 4.4, we have $\alpha_2 + \delta_{1j} \in R^W$ for $j = 1, \cdots, b$. For arbitrary $\gamma \in S_5'$, we get $s_{\alpha_1}s_{\alpha_{1j}}s_{\alpha_2}s_{\alpha_2+\delta_{1j}}(\gamma) = \gamma + \delta_{1j}$ and $s_{\alpha_2}s_{\alpha_{2j}}(\gamma) = \gamma + 3\delta_{2j}$. Therefore, if $\gamma \in R^W$, we obtain $\gamma \pm \delta_{1j}, \gamma \pm 3\delta_{2k} \in R^W$ for all $j = 1, \cdots, b$ and $k = 1, \cdots, s$. By induction on the number $n = \sum\limits_{j=1}^{b} |a_j| + 3\sum\limits_{j=1}^{s} |b_j|$, we have $S_5' \subseteq R_l^W$.

(3) Consider the type $A_1^{(b,0,t)}$. Let

$$\begin{aligned} S_6 : \ = \ & \{\pm\alpha_1 + \sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{t} b_j\delta_{1j}' \mid a_j, b_j \in \mathbf{Z}, \\ & \text{and there is at most one odd } a_j, j = 1 \cdots, b\}. \end{aligned}$$

Since $\alpha_{1j} = \alpha_1 + \delta_{1j}, j = 1, \cdots, b$, we have $\Delta_s \subseteq S_6$. And for any $\beta = \alpha + \sum\limits_{j=1}^{b} a_j\delta_{1j} + \sum\limits_{j=1}^{t} b_j\delta_{1j}' \in S_6$ with $\alpha = \pm\alpha_1$, we obtain

$$s_{\alpha_{1k}}(\beta) = s_{\alpha_{1k}}(\alpha) + \sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{t} b_j\delta_{1j}' = s_{\alpha_1}(\alpha) - (\alpha_1^\vee, \alpha)\delta_{1k} + \sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{t} b_j\delta_{1j}';$$

$$s_{\beta_{1k}}(\beta) = s_{\beta_{1k}}(\alpha) + \sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{t} b_j\delta_{1j}' = s_{\alpha_1}(\alpha) - \frac{1}{2}(\alpha_1^\vee, \alpha)\delta_{1k}' + \sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{t} b_j\delta_{1j}'.$$

Note that $(\alpha_1^\vee, \alpha) = \pm 2$. Then we get

$$s_{\alpha_{11}}(\beta), \cdots, s_{\alpha_{1b}}(\beta), s_{\beta_{11}'}(\beta), \cdots, s_{\beta_{1t}'}(\beta) \in S_6,$$

and $s_{\alpha_1}(\beta) = s_{\alpha_1}(\alpha) + \sum\limits_{j=1}^{b} a_j\delta_{1j} + \sum\limits_{j=1}^{t} b_j\delta_{1j}' \in S_6$. So by induction on the number of reflections, we obtain $R_s^W \subseteq S_6$.

It remains to show $S_6 \subseteq R_s^W$. We only need to show $S_6' \subseteq R_s^W$, where $S_6' := \{\alpha_1 + \sum\limits_{j=1}^{b} a_j\delta_{1j} + \sum\limits_{j=1}^{t} b_j\delta_{1j}' \mid a_j, b_j \in \mathbf{Z}$, and there is at most one odd $a_j, j = 1 \cdots, b\}$. For arbitrary $\gamma \in S_6'$, we obtain $s_{\alpha_{1j}}s_{\alpha_1}(\gamma) = \gamma + 2\delta_{1j}$ and $s_{\beta_{1j}}s_{\alpha_1}(\gamma) = \gamma + \delta_{1j}'$. Therefore, if $\gamma \in R^W$, we have $\gamma \pm 2\delta_{1j}, \gamma \pm \delta_{1k}' \in R^W$ for all $j = 1, \cdots, b$ and $k = 1, \cdots, s$. Moreover, $\alpha_1 + \delta_{1j} = \alpha_{1j} \in R_s^W$ for $j = 1, \cdots, b$. By induction on the number $n = \sum\limits_{j=1}^{b} |a_j| + \sum\limits_{j=1}^{t} |b_j|$, we get $S_6' \subseteq R_s^W$.

So $R_s^W = S_6 = \{\pm\alpha_1 + \sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{t} b_j\delta'_{1j} \mid a_j, b_j \in \mathbf{Z},$ and there is at most one odd $a_j, j = 1 \cdots, b\}$.

Let $S_7 := \{\pm 2\alpha_1 + 4\sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{t} b_j\delta'_{1j} \mid a_j, b_j \in \mathbf{Z},$ and there is exactly one odd $b_j, j = 1, \cdots, t\}$. Since $\beta_{1j} = 2\alpha_1 + \delta'_{1j}, j = 1, \cdots, t$, we have $\Delta_l \subseteq S_7$. And for any $\beta = \alpha + 4\sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{t} b_j\delta'_{1j} \in S_7$ with $\alpha = \pm 2\alpha_1$, we have

$$s_{\alpha_{1k}}(\beta) = s_{\alpha_{1k}}(\alpha) + 4\sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{t} b_j\delta'_{1j} = s_{\alpha_1}(\alpha) - (\alpha_1^\vee, \alpha)\delta_{1k} + 4\sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{t} b_j\delta'_{1j};$$

$$s_{\beta_{1k}}(\beta) = s_{\beta_{1k}}(\alpha) + 4\sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{t} b_j\delta'_{1j} = s_{\alpha_1}(\alpha) - \frac{1}{2}(\alpha_1^\vee, \alpha)\delta'_{1k} + 4\sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{t} b_j\delta'_{1j}.$$

Moreover, $s_{\alpha_1}(\alpha) = -\alpha,\ (\alpha_1^\vee, \alpha) = \pm 4$. Then we obtain

$$s_{\alpha_{11}}(\beta), \cdots, s_{\alpha_{1b}}(\beta), s_{\beta_{11}}(\beta), \cdots, s_{\beta_{1t}}(\beta) \in S_7,$$

and $s_{\alpha_1}(\beta) = s_{\alpha_1}(\alpha) + 4\sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{t} b_j\delta'_{1j} \in S_7$. So by induction on the number of reflections, we get $R_l^W \subseteq S_7$.

Next we prove $S_7 \subseteq R_l^W$. It suffices to show $S'_7 \subseteq R_l^W$, where $S'_7 := \{2\alpha_1 + 4\sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{t} b_j\delta'_{1j} \mid a_j, b_j \in \mathbf{Z},$ and there is exactly one odd $b_j, j = 1, \cdots, t\}$. For any $\gamma \in S'_7$, we get $s_{\alpha_{1j}}s_{\alpha_1}(\gamma) = \gamma + 4\delta_{1j}$ and $s_{\beta_{1j}}s_{\alpha_1}(\gamma) = \gamma + 2\delta'_{1j}$. So, if $\gamma \in R^W$, then $\gamma \pm 4\delta_{1j}, \gamma \pm 2\delta'_{1k} \in R^W$ for all $j = 1, \cdots, b$ and $k = 1, \cdots, s$. Notice that $2\alpha_1 + \delta'_{1j} = \beta_{1j} \in R_l^W$ for $j = 1, \cdots, t$. So by induction on the number $n = 4\sum_{j=1}^{b} |a_j| + \sum_{j=1}^{t} |b_j|$, we obtain that $S'_7 \subseteq R_l^W$.

Hence $R_l^W = S_7 = \{\pm 2\alpha_1 + 4\sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{t} b_j\delta'_{1j} \mid a_j, b_j \in \mathbf{Z},$ and there is exactly one odd $b_j, j = 1, \cdots, t\}$.

(4) Consider the type $B_l^{(b,s,t)}$. Let $S_8 := \{\alpha + \sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{s} b_j\delta_{lj} + \sum_{j=1}^{t} c_j\delta'_{lj} \mid \alpha \in \dot{R}_s, a_j, b_j, c_j \in \mathbf{Z},$ and there is at most one odd $b_j, j = 1, \cdots, s\}$. Since $\alpha_{lj} = \alpha_l + \delta_{lj}, j = 1, \cdots, s$, we have $\Delta_s \subseteq S_8$. And for any $\beta = \alpha + \sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{s} b_j\delta_{lj} + \sum_{j=1}^{t} c_j\delta'_{lj} \in S_8$ with $\alpha \in \dot{R}_s$, we obtain

$$\begin{aligned} s_{\alpha_{1k}}(\beta) &= s_{\alpha_{1k}}(\alpha) + \sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{s} b_j\delta_{lj} + \sum_{j=1}^{t} c_j\delta'_{lj} \\ &= s_{\alpha_1}(\alpha) - (\alpha_1^\vee, \alpha)\delta_{1k} + \sum_{j=1}^{b} a_j\delta_{1j} + \sum_{j=1}^{s} b_j\delta_{lj} + \sum_{j=1}^{t} c_j\delta'_{lj}; \end{aligned}$$

$$\begin{aligned} s_{\alpha_{lk}}(\beta) &= s_{\alpha_{lk}}(\alpha)+\sum_{j=1}^{b}a_j\delta_{1j}+\sum_{j=1}^{s}b_j\delta_{lj}+\sum_{j=1}^{t}c_j\delta'_{lj} \\ &= s_{\alpha_l}(\alpha)-(\alpha_l^\vee,\alpha)\delta_{lk}+\sum_{j=1}^{b}a_j\delta_{1j}+\sum_{j=1}^{s}b_j\delta_{lj}+\sum_{j=1}^{t}c_j\delta'_{lj}; \end{aligned}$$

$$\begin{aligned} s_{\beta_{lk}}(\beta) &= s_{\beta_{lk}}(\alpha)+\sum_{j=1}^{b}a_j\delta_{1j}+\sum_{j=1}^{s}b_j\delta_{lj}+\sum_{j=1}^{t}c_j\delta'_{lj} \\ &= s_{\alpha_l}(\alpha)-\frac{1}{2}(\alpha_l^\vee,\alpha)\delta'_{lk}+\sum_{j=1}^{b}a_j\delta_{1j}+\sum_{j=1}^{s}b_j\delta_{lj}+\sum_{j=1}^{t}c_j\delta'_{lj}. \end{aligned}$$

By Lemma 4.5, $(\alpha_l^\vee,\alpha)$ is even. Thus we have

$$s_{\alpha_{11}}(\beta),\cdots,s_{\alpha_{1b}}(\beta),s_{\alpha_{l1}}(\beta),\cdots,s_{\alpha_{ls}}(\beta),s_{\beta_{l1}}(\beta),\cdots,s_{\beta_{lt}}(\beta)\in S_8,$$

and $s_{\alpha_i}(\beta)=s_{\alpha_i}(\alpha)+\sum\limits_{j=1}^{b}a_j\delta_{1j}+\sum\limits_{j=1}^{s}b_j\delta_{lj}+\sum\limits_{j=1}^{t}c_j\delta'_{lj}\in S_8$ for $i=1,\cdots,l$. So by induction on the number of reflections, we get $R_s^W\subseteq S_8$.

It remains to prove $S_8\subseteq R_s^W$. We only need to show that $S'_8\subseteq R_s^W$, where $S'_8:=\{\alpha_l+\sum\limits_{j=1}^{b}a_j\delta_{1j}+\sum\limits_{j=1}^{s}b_j\delta_{lj}+\sum\limits_{j=1}^{t}c_j\delta'_{lj}\mid a_j,b_j,c_j\in\mathbf{Z},$ and there is at most one odd b_j, $j=1,\cdots,s\}$. By Lemma 4.4, we have $\alpha_i+\delta_{1j}\in R^W$ for $i=1,\cdots,l,j=1,\cdots,b$. For any $\gamma\in S'_8$, we get $s_{\alpha_{l-1}}s_{\alpha_{l-1}+\delta_{1j}}(\gamma)=\gamma+\delta_{1j}$, $s_{\alpha_{lj}}s_{\alpha_l}(\gamma)=\gamma+2\delta_{lj}$ and $s_{\beta_{lj}}s_{\alpha_l}(\gamma)=\gamma+\delta'_{lj}$. Therefore, if $\gamma\in R^W$, we have $\gamma\pm\delta_{1i},\gamma\pm2\delta_{lj},\gamma\pm\delta'_{lk}\in R^W$ for all $i=1,\cdots,b,j=1,\cdots,s$ and $k=1,\cdots,t$. Note that $\alpha_l+\delta_{lj}=\alpha_{lj}\in R_s^W$ for $j=1,\cdots,s$. By induction on the number $n=\sum\limits_{j=1}^{b}|a_j|+\sum\limits_{j=1}^{s}|b_j|+\sum\limits_{j=1}^{t}|c_j|$, we obtain $S'_8\subseteq R_s^W$.

So $R_s^W=S_8=\{\alpha+\sum\limits_{j=1}^{b}a_j\delta_{1j}+\sum\limits_{j=1}^{s}b_j\delta_{lj}+\sum\limits_{j=1}^{t}c_j\delta'_{lj}\mid\alpha\in\dot{R}_s,a_j,b_j,c_j\in\mathbf{Z},$ and there is at most one odd $b_j,j=1,\cdots,s\}$.

Let $S_9:=\{2\alpha+2\sum\limits_{j=1}^{b}a_j\delta_{1j}+4\sum\limits_{j=1}^{s}b_j\delta_{lj}+\sum\limits_{j=1}^{t}c_j\delta'_{lj}\mid\alpha\in\dot{R}_s,a_j,b_j,c_j\in\mathbf{Z},$ and there is exactly one odd $c_j,j=1,\cdots,t\}$. Since $\beta_{lj}=2\alpha_l+\delta'_{lj},j=1,\cdots,t$, we get $\Delta_l\subseteq S_9$. And for any $\beta=2\alpha+2\sum\limits_{j=1}^{b}a_j\delta_{1j}+4\sum\limits_{j=1}^{s}b_j\delta_{lj}+\sum\limits_{j=1}^{t}c_j\delta'_{lj}\in S_9$ with $\alpha\in\dot{R}_s$, we have

$$\begin{aligned} s_{\alpha_{1k}}(\beta) &= 2s_{\alpha_{1k}}(\alpha)+2\sum_{j=1}^{b}a_j\delta_{1j}+4\sum_{j=1}^{s}b_j\delta_{lj}+\sum_{j=1}^{t}c_j\delta'_{lj} \\ &= 2s_{\alpha_1}(\alpha)-2(\alpha_1^\vee,\alpha)\delta_{1k}+2\sum_{j=1}^{b}a_j\delta_{1j}+4\sum_{j=1}^{s}b_j\delta_{lj}+\sum_{j=1}^{t}c_j\delta'_{lj}; \end{aligned}$$

$$\begin{aligned} s_{\alpha_{lk}}(\beta) &= 2s_{\alpha_{lk}}(\alpha) + 2\sum_{j=1}^{b} a_j\delta_{1j} + 4\sum_{j=1}^{s} b_j\delta_{lj} + \sum_{j=1}^{t} c_j\delta'_{lj} \\ &= 2s_{\alpha_l}(\alpha) - 2(\alpha_l^\vee, \alpha)\delta_{lk} + 2\sum_{j=1}^{b} a_j\delta_{1j} + 4\sum_{j=1}^{s} b_j\delta_{lj} + \sum_{j=1}^{t} c_j\delta'_{lj}; \end{aligned}$$

$$\begin{aligned} s_{\beta_{lk}}(\beta) &= 2s_{\beta_{lk}}(\alpha) + 2\sum_{j=1}^{b} a_j\delta_{1j} + 4\sum_{j=1}^{s} b_j\delta_{lj} + \sum_{j=1}^{t} c_j\delta'_{lj} \\ &= 2s_{\alpha_l}(\alpha) - (\alpha_l^\vee, \alpha)\delta'_{lk} + 2\sum_{j=1}^{b} a_j\delta_{1j} + 4\sum_{j=1}^{s} b_j\delta_{lj} + \sum_{j=1}^{t} c_j\delta'_{lj}. \end{aligned}$$

By Lemma 4.5, $(\alpha_l^\vee, \alpha)$ is even. Then we obtain

$$s_{\alpha_{11}}(\beta), \cdots, s_{\alpha_{1b}}(\beta), s_{\alpha_{l1}}(\beta), \cdots, s_{\alpha_{ls}}(\beta), s_{\beta_{l1}}(\beta), \cdots, s_{\beta_{lt}}(\beta) \in S_9,$$

and $s_{\alpha_i}(\beta) = 2s_{\alpha_i}(\alpha) + 2\sum\limits_{j=1}^{b} a_j\delta_{1j} + 4\sum\limits_{j=1}^{s} b_j\delta_{lj} + \sum\limits_{j=1}^{t} c_j\delta'_{lj} \in S_9$ for $i = 1, \cdots, l$. So by induction on the number of reflections, we obtain $R_l^W \subseteq S_9$.

Next we prove $S_9 \subseteq R_l^W$. It suffices to show that $S'_9 \subseteq R_l^W$, where $S'_9 := \{2\alpha_l + 2\sum\limits_{j=1}^{b} a_j\delta_{1j} + 4\sum\limits_{j=1}^{s} b_j\delta_{lj} + \sum\limits_{j=1}^{t} c_j\delta'_{lj} \mid a_j, b_j, c_j \in \mathbf{Z},$ and there is exactly one odd $c_j, j = 1, \cdots, t\}$. By Lemma 4.4, we have $\alpha_i + \delta_{1j} \in R^W$ for $i = 1, \cdots, l$ and $j = 1, \cdots, b$. For any $\gamma \in S'_9$, we get $s_{\alpha_{l-1}} s_{\alpha_{l-1}+\delta_{1j}}(\gamma) = \gamma + 2\delta_{1j}$, $s_{\alpha_{lj}} s_{\alpha_l}(\gamma) = \gamma + 4\delta_{lj}$ and $s_{\beta_{lj}} s_{\alpha_l}(\gamma) = \gamma + 2\delta'_{lj}$. Therefore, if $\gamma \in R^W$, we obtain $\gamma \pm 2\delta_{1i}, \gamma \pm 4\delta_{lj}, \gamma \pm 2\delta'_{lk} \in R^W$ for all $i = 1, \cdots, b, j = 1, \cdots, s$ and $k = 1, \cdots, t$. Note that $2\alpha_l + \delta'_{lj} = \beta_{lj} \in R_l^W$ for $j = 1, \cdots, s$. By induction on the number $n = 2\sum\limits_{j=1}^{b} |a_j| + 4\sum\limits_{j=1}^{s} |b_j| + \sum\limits_{j=1}^{t} |c_j|$, we have $S'_9 \subseteq R_l^W$.

So $R_l^W = S_9 = \{2\alpha + 2\sum\limits_{j=1}^{b} a_j\delta_{1j} + 4\sum\limits_{j=1}^{s} b_j\delta_{lj} + \sum\limits_{j=1}^{t} c_j\delta'_{lj} \mid \alpha \in \dot{R}_s, a_j, b_j, c_j \in \mathbf{Z},$ and there is exactly one odd $c_j, j = 1, \cdots, t\}$.

(4.1) $X_l = B_l (l = 2)$. Let S_{10} be the set on the right-hand side. Since $\alpha_{1j} = \alpha_1 + \delta_{1j}, j = 1, \cdots, b$, we have $\Delta_m \subseteq S_{10}$. And for any $\beta = \alpha + \sum\limits_{j=1}^{b} a_j\delta_{1j} + 2\sum\limits_{j=1}^{s} b_j\delta_{2j} + \sum\limits_{j=1}^{t} c_j\delta'_{2j} \in S_{10}$ with $\alpha \in \dot{R}_l$, we have

$$\begin{aligned} s_{\alpha_{1k}}(\beta) &= s_{\alpha_{1k}}(\alpha) + \sum_{j=1}^{b} a_j\delta_{1j} + 2\sum_{j=1}^{s} b_j\delta_{2j} + \sum_{j=1}^{t} c_j\delta'_{2j} \\ &= s_{\alpha_1}(\alpha) - (\alpha_1^\vee, \alpha)\delta_{1k} + \sum_{j=1}^{b} a_j\delta_{1j} + 2\sum_{j=1}^{s} b_j\delta_{2j} + \sum_{j=1}^{t} c_j\delta'_{2j}; \end{aligned}$$

$$\begin{aligned} s_{\alpha_{2k}}(\beta) &= s_{\alpha_{2k}}(\alpha)+\sum_{j=1}^{b}a_j\delta_{1j}+2\sum_{j=1}^{s}b_j\delta_{2j}+\sum_{j=1}^{t}c_j\delta'_{2j} \\ &= s_{\alpha_2}(\alpha)-(\alpha_2^\vee,\alpha)\delta_{2k}+\sum_{j=1}^{b}a_j\delta_{1j}+2\sum_{j=1}^{s}b_j\delta_{2j}+\sum_{j=1}^{t}c_j\delta'_{2j}; \end{aligned}$$

$$\begin{aligned} s_{\beta_{2k}}(\beta) &= s_{\beta_{2k}}(\alpha)+\sum_{j=1}^{b}a_j\delta_{1j}+2\sum_{j=1}^{s}b_j\delta_{2j}+\sum_{j=1}^{t}c_j\delta'_{2j} \\ &= s_{\alpha_2}(\alpha)-\frac{1}{2}(\alpha_2^\vee,\alpha)\delta'_{2k}+\sum_{j=1}^{b}a_j\delta_{1j}+2\sum_{j=1}^{s}b_j\delta_{2j}+\sum_{j=1}^{t}c_j\delta'_{2j}. \end{aligned}$$

By Lemma 4.5, $(\alpha_i^\vee,\alpha)$ is even for $i=1,2$. Then we have

$$s_{\alpha_{11}}(\beta),\cdots,s_{\alpha_{1b}}(\beta),s_{\alpha_{21}}(\beta),\cdots,s_{\alpha_{2s}}(\beta),s_{\beta_{21}}(\beta),\cdots,s_{\beta_{2t}}(\beta)\in S_{10},$$

and $s_{\alpha_i}(\beta)=s_{\alpha_i}(\alpha)+\sum\limits_{j=1}^{b}a_j\delta_{1j}+2\sum\limits_{j=1}^{s}b_j\delta_{2j}+\sum\limits_{j=1}^{t}c_j\delta'_{2j}\in S_{10}$ for $i=1,2$. So by induction on the number of reflections, we obtain $R_m^W\subseteq S_{10}$.

Next we prove $S_{10}\subseteq R_m^W$. It suffices to show that $S'_{10}\subseteq R_m^W$, where $S'_{10}:=\{\alpha_1+\sum\limits_{j=1}^{b}a_j\delta_{1j}+2\sum\limits_{j=1}^{s}b_j\delta_{2j}+\sum\limits_{j=1}^{t}c_j\delta'_{2j}\mid a_j,b_j,c_j\in\mathbf{Z},$ and there is at most one odd $a_j, j=1\cdots,b\}$. For any $\gamma\in S'_{10}$, we get $s_{\alpha_{1j}}s_{\alpha_1}(\gamma)=\gamma+2\delta_{1j}$, $s_{\alpha_2}s_{\alpha_{2j}}(\gamma)=\gamma+2\delta_{2j}$ and $s_{\alpha_2}s_{\beta_{2j}}(\gamma)=\gamma+\delta'_{2j}$. Therefore, if $\gamma\in R^W$, we have $\gamma\pm2\delta_{1i},\gamma\pm2\delta_{2j},\gamma\pm\delta'_{2k}\in R^W$ for all $i=1,\cdots,b, j=1,\cdots,s$ and $k=1,\cdots,t$. Note that $\alpha_1+\delta_{1j}=\alpha_{1j}\in R_m^W$ for $j=1,\cdots,b$. Hence, by induction on the number $n=\sum\limits_{j=1}^{b}|a_j|+2\sum\limits_{j=1}^{s}|b_j|+\sum\limits_{j=1}^{t}|c_j|$, we obtain $S'_{10}\subseteq R_m^W$.

(4.2) $X_l=B_l(l>2)$. Let S_{11} be the set on the right-hand side. Since $\alpha_{1j}=\alpha_1+\delta_{1j}, j=1,\cdots,b$, we have $\Delta_m\subseteq S_{11}$. And for any $\beta=\alpha+\sum\limits_{j=1}^{b}a_j\delta_{1j}+2\sum\limits_{j=1}^{s}b_j\delta_{lj}+\sum\limits_{j=1}^{t}c_j\delta'_{lj}\in S_{11}$ with $\alpha\in\dot{R}_l$, we have

$$\begin{aligned} s_{\alpha_{1k}}(\beta) &= s_{\alpha_{1k}}(\alpha)+\sum_{j=1}^{b}a_j\delta_{1j}+2\sum_{j=1}^{s}b_j\delta_{lj}+\sum_{j=1}^{t}c_j\delta'_{lj} \\ &= s_{\alpha_1}(\alpha)-(\alpha_1^\vee,\alpha)\delta_{1k}+\sum_{j=1}^{b}a_j\delta_{1j}+2\sum_{j=1}^{s}b_j\delta_{lj}+\sum_{j=1}^{t}c_j\delta'_{lj}; \end{aligned}$$

$$\begin{aligned} s_{\alpha_{lk}}(\beta) &= s_{\alpha_{lk}}(\alpha)+\sum_{j=1}^{b}a_j\delta_{1j}+2\sum_{j=1}^{s}b_j\delta_{lj}+\sum_{j=1}^{t}c_j\delta'_{lj} \\ &= s_{\alpha_l}(\alpha)-(\alpha_l^\vee,\alpha)\delta_{lk}+\sum_{j=1}^{b}a_j\delta_{1j}+2\sum_{j=1}^{s}b_j\delta_{lj}+\sum_{j=1}^{t}c_j\delta'_{lj}; \end{aligned}$$

$$\begin{aligned} s_{\beta_{lk}}(\beta) &= s_{\beta_{lk}}(\alpha)+\sum_{j=1}^{b}a_j\delta_{1j}+2\sum_{j=1}^{s}b_j\delta_{lj}+\sum_{j=1}^{t}c_j\delta'_{lj}\\ &= s_{\alpha_l}(\alpha)-\frac{1}{2}(\alpha_l^\vee,\alpha)\delta'_{lk}+\sum_{j=1}^{b}a_j\delta_{1j}+2\sum_{j=1}^{s}b_j\delta_{lj}+\sum_{j=1}^{t}c_j\delta'_{lj}. \end{aligned}$$

By Lemma 4.5, $(\alpha_l^\vee,\alpha)$ is even. Then we obtain

$$s_{\alpha_{11}}(\beta),\cdots,s_{\alpha_{1b}}(\beta),s_{\alpha_{l1}}(\beta),\cdots,s_{\alpha_{ls}}(\beta),s_{\beta_{l1}}(\beta),\cdots,s_{\beta_{lt}}(\beta)\in S_{11},$$

and $s_{\alpha_i}(\beta)=s_{\alpha_i}(\alpha)+\sum\limits_{j=1}^{b}a_j\delta_{1j}+2\sum\limits_{j=1}^{s}b_j\delta_{lj}+\sum\limits_{j=1}^{t}c_j\delta'_{lj}\in S_{11}$ for $i=1,\cdots,l$. So by induction on the number of reflections, we obtain $R_m^W\subseteq S_{11}$.

It remains to prove $S_{11}\subseteq R_m^W$. By Lemma 4.4, we only need to show $S'_{11}\subseteq R_m^W$, where $S'_{11}:=\{\alpha_{l-1}+\sum\limits_{j=1}^{b}a_j\delta_{1j}+2\sum\limits_{j=1}^{s}b_j\delta_{lj}+\sum\limits_{j=1}^{t}c_j\delta'_{lj}\mid a_j,b_j,c_j\in\mathbf{Z}\}$. By Lemma 4.4, we have $\alpha_i+\delta_{1j}\in R^W$ for $i=1,\cdots,l,j=1,\cdots,b$. For any $\gamma\in S'_{11}$, we obtain $s_{\alpha_{l-2}}s_{\alpha_{l-2}+\delta_{1j}}(\gamma)=\gamma+\delta_{1j}$, $s_{\alpha_l}s_{\alpha_{lj}}(\gamma)=\gamma+2\delta_{lj}$ and $s_{\alpha_l}s_{\beta_{lj}}(\gamma)=\gamma+\delta'_{lj}$. Therefore, if $\gamma\in R^W$, we have $\gamma\pm\delta_{1i},\gamma\pm2\delta_{lj},\gamma\pm\delta'_{lk}\in R^W$ for all $i=1,\cdots,b,j=1,\cdots,s$ and $k=1,\cdots,t$. By induction on the number $n=\sum\limits_{j=1}^{b}|a_j|+2\sum\limits_{j=1}^{s}|b_j|+\sum\limits_{j=1}^{t}|c_j|$, we obtain $S'_{11}\subseteq R_m^W$. □

References

[AABGP] B. Allison, N. Azam, S. Berman, Y. Gao, A. Pianzola, Extended affine Lie algebras and their root systems, Memoir. A. M. S., 126 (1997).

[APR] M. Auslander, M. I. Platzeck, I. Reiten, Coxeter functors without diagrams, Trans. AMS 250 (1979), 1-46.

[BM] S. Berman, R. V. Moody, Lie algebras graded by finite root systems and the intersection matrix algebras of Slodowy, Invent. math. 108 (1992), 323-347.

[BZ] G. Benkart, E. Zelmanov, Lie algebras graded by finite root systems and intersection matrix algebras, Invent. Math. 126 (1996), 1-45.

[Kac] V. G. Kac, Infinite dimensional Lie algebras, third edition, Cambridge University press, 1990.

[Peng] L. Peng, Intersection matrix Lie algebras and Ringel-Hall Lie algebras of tilted algebras, Proc. the 9th Inter. Conf. on Representations of Algebras, (1)(2002), 98-108,

[Sai] K. Saito, Extended Affine Root system I (Coxeter Transformations), Publ. RIMS. Kyoto Univ., 21 (1985), 75-179.

[Slo1] P. Slodowy, Singularitäten, Kac-Moody-Liealgebren, assoziierte Gruppen und Verallgemeinerungen, Habiliationsschrift, Universität Bonn, March 1984.

[Slo2] P. Slodowy, Beyond Kac-Moody algebras, and inside, Canad. Math. Soc. Conf. Proc., 5 (1988), 361-371.

[SY] K. Saito, D. Yoshii, Extended affine root system IV (Simple laced ellipic Lie algebras), Publ. RIMS. Kyoto Univ., 36 (2000), 385-421.

[XP] M. Xu, L. G. Peng, Symmetrizable Intersection Matrix Lie Algebras, Algebra Colloquium (to appear).

Department of Mathematics, Sichuan University, 610064 Chengdu, China
E-mail address: `penglg@scu.edu.cn`

Department of Mathematics, Southwest Jiaotong University, 610031 chengdu, China
E-mail address: `xumang@home.swjtu.edu.cn`